CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 100, 2007

A

Abel, C., 759
Adamczyk, J. J., Jr., 1569
Adan, A., 1789
Addison, S., 1844
Addo-Bediako, A., 315
Ahmad, A., 1221
Ahmad, M., 886
Aiken, R. M., 1248, 1555
Alatawi, F. J., 1904
Aldrich, J. R., 680, 1751
Ali, M. I., 921
Allen, D. C., 335
Alm, S. R., 1604
Alma, A., 1910
Alphey, L., 1642
Alphey, N., 1642
Altosaar, I., 172
Aluko, G. A., 1037
Alvarez, J. M., 646
Álvarez-Alfageme, F., 195
Alyokhin, A., 1871
Anderbrant, O., 1017
Andow, D. A., 164, 195
Andreadis, S. S., 195
Antwi, F. B., 710, 1201
Appel, A. G., 871
Arango, R., 172
Aref, S., 1241
Arévalo, H. A., 1622
Armstrong, J. W., 1091
Arredondo, J., 1153
Arthur, F. H., 604, 1714
Arthurs, S. P., 702
Aspaly, G., 586
Athanassiou, C. G., 599, 1724
Atterholt, C. A., 1316
Azuma, M., 1276

B

Bagwell, R., 1916
Bailer, A. J., 1945
Baker, C., 1679
Baker, C. A., 1685
Baker, G. T., 976
Baker, M. B., 1871
Baker, P. B., 1381
Baniameri, V., 350
Banks, J. E., 1027
Baronio, P., 26
Barreda-Landa, A., 752
Barrera, J. F., 291
Barrett, B. A., 72
Barros, W., 551
Bartlett, J. S., 1136
Bashasab, F., 903
Baumler, R. E., 451
Baxendale, F., 136
Beaman, L. D., 267
Beiriger, R., 1887
Bell, V., 745
Bellec, C., 545
Beres, B. L., 79
Bergh, J. C., 405
Beukeboom, L. W., 1033
Bianco, P., 1047
Bielza, P., 916
Bikoba, V., 1084
Bishop, B., 307
Bixby, A., 1604
Bland, J., 880
Bohac, J. R., 566
Boll, R., 225
Bommireddy, P. L., 1560
Bonning, B., 1862
Bonsi, C. K., 315
Boonsuepsakul, S., 946
Boopathy, R., 880
Booth, L. H., 11
Bosco, D., 1504
Bostanian, N. J., 1210
Bosu, P. P., 1808
Botha, A.-M., 990
Boukouvala, M. C., 599
Bowling, B., 1679
Bowling, R. D., 1221
Bradley, J. R., Jr., 180, 801
Bradshaw, J. D., 808
Brandenburg, R. L., 1241
Bridges, W. C., Jr., 1229
Brodeur, J., 1344
Brooks, T. D., 1470
Brosius, T. R., 1268
Brown, M. D., 61
Brown, S. L., 1704
Buckley, P. M., 1470
Budia, F., 773, 1789
Buntin, G. D., 1704
Burd, J. D., 1679, 1692
Burgio, G., 298
Burkness, E. C., 1000
Burris, E., 1916
Buschman, L. L., 1862
Bushman, B. S., 1470
Buss, G. R., 962, 1456
Butler, S. M., 1489
Butrón, A., 209
Büyükgüzel, E., 1533
Byers, J. R., 79

C

Callcott, A.-M., 307
Calvin, D. D., 1129
Camelo, L. D. A., 366
Campbell, J. F., 1714
Campbell, L. R., 984, 1657
Campos, S. E., 291
Cánovas, F., 123
Cao, M. Z., 1854
Capinera, J. L., 117
Cappaert, D., 1304
Cárcamo, H. A., 79
Cardona, C., 1896
Carey, D. R., 1201
Carlson, D. A., 483, 613
Carlton, C. E., 54
Caroli, L., 1637
Carrière, Y., 1650
Carroll, J., 1400
Carstens, J., 136
Casada, M. E., 604
Casana-Giner, V., 1124
Casati, P., 1047
Castánera, P., 195
Castañera, P., 1444, 1836
Castle, S. J., 1053
Castrejón-Gómez, V. R., 1173
Castro, A. M., 1781
Catchot, A., 1916
Chandler, L. D., 1542
Chaney, W. E., 39
Chang, C. L., 322
Chapin, J. W., 1229
Chareonviriyaphap, T., 545
Charlet, L. D., 1248, 1555
Chaudhury, M. F., 241
Chauhan, K. R., 1751
Chen, Q., 1450
Chen, W.-L., 685
Chen, W. M., 1854
Chen, Y., 1821
Cherry, S., 381
Chiasson, H., 459
Chicca, M., 1637
Chilcutt, C. F., 88, 327, 1428
Childers, C. C., 233
Choo, H. Y., 103
Chouinard, G., 737
Cinanni, L., 1047
Civolani, S., 1637
Clark, E. L., 1304
Clark, T. L., 557
Clarke, S. R., 1298
Clement, S. L., 131
Cloyd, R. A., 459, 1353

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 100, 2007

A

Abel, C., 759
Adamczyk, J. J., Jr., 1569
Adan, A., 1789
Addison, S., 1844
Addo-Bediako, A., 315
Ahmad, A., 1221
Ahmad, M., 886
Aiken, R. M., 1248, 1555
Alatawi, F. J., 1904
Aldrich, J. R., 680, 1751
Ali, M. I., 921
Allen, D. C., 335
Alm, S. R., 1604
Alma, A., 1910
Alphey, L., 1642
Alphey, N., 1642
Altosaar, I., 172
Aluko, G. A., 1037
Alvarez, J. M., 646
Álvarez-Alfageme, F., 195
Alyokhin, A., 1871
Anderbrant, O., 1017
Andow, D. A., 164, 195
Andreadis, S. S., 195
Antwi, F. B., 710, 1201
Appel, A. G., 871
Arango, R., 172
Aref, S., 1241
Arévalo, H. A., 1622
Armstrong, J. W., 1091
Arredondo, J., 1153
Arthur, F. H., 604, 1714
Arthurs, S. P., 702
Aspaly, G., 586
Athanassiou, C. G., 599, 1724
Atterholt, C. A., 1316
Azuma, M., 1276

B

Bagwell, R., 1916
Bailer, A. J., 1945
Baker, C., 1679
Baker, C. A., 1685
Baker, G. T., 976
Baker, M. B., 1871
Baker, P. B., 1381
Baniameri, V., 350
Banks, J. E., 1027
Baronio, P., 26
Barreda-Landa, A., 752
Barrera, J. F., 291
Barrett, B. A., 72
Barros, W., 551
Bartlett, J. S., 1136
Bashasab, F., 903
Baumler, R. E., 451
Baxendale, F., 136
Beaman, L. D., 267
Beiriger, R., 1887
Bell, V., 745
Bellec, C., 545
Beres, B. L., 79
Bergh, J. C., 405
Beukeboom, L. W., 1033
Bianco, P., 1047
Bielza, P., 916
Bikoba, V., 1084
Bishop, B., 307
Bixby, A., 1604
Bland, J., 880
Bohac, J. R., 566
Boll, R., 225
Bommireddy, P. L., 1560
Bonning, B., 1862
Bonsi, C. K., 315
Boonsuepsakul, S., 946
Boopathy, R., 880
Booth, L. H., 11
Bosco, D., 1504
Bostanian, N. J., 1210
Bosu, P. P., 1808
Botha, A.-M., 990
Boukouvala, M. C., 599
Bowling, B., 1679
Bowling, R. D., 1221
Bradley, J. R., Jr., 180, 801
Bradshaw, J. D., 808
Brandenburg, R. L., 1241
Bridges, W. C., Jr., 1229
Brodeur, J., 1344
Brooks, T. D., 1470
Brosius, T. R., 1268
Brown, M. D., 61
Brown, S. L., 1704
Buckley, P. M., 1470
Budia, F., 773, 1789
Buntin, G. D., 1704
Burd, J. D., 1679, 1692
Burgio, G., 298
Burkness, E. C., 1000
Burris, E., 1916
Buschman, L. L., 1862
Bushman, B. S., 1470
Buss, G. R., 962, 1456
Butler, S. M., 1489
Butrón, A., 209
Büyükgüzel, E., 1533
Byers, J. R., 79

C

Callcott, A.-M., 307
Calvin, D. D., 1129
Camelo, L. D. A., 366
Campbell, J. F., 1714
Campbell, L. R., 984, 1657
Campos, S. E., 291
Cánovas, F., 123
Cao, M. Z., 1854
Capinera, J. L., 117
Cappaert, D., 1304
Cárcamo, H. A., 79
Cardona, C., 1896
Carey, D. R., 1201
Carlson, D. A., 483, 613
Carlton, C. E., 54
Caroli, L., 1637
Carrière, Y., 1650
Carroll, J., 1400
Carstens, J., 136
Casada, M. E., 604
Casana-Giner, V., 1124
Casati, P., 1047
Castánera, P., 195
Castañera, P., 1444, 1836
Castle, S. J., 1053
Castrejón-Gómez, V. R., 1173
Castro, A. M., 1781
Catchot, A., 1916
Chandler, L. D., 1542
Chaney, W. E., 39
Chang, C. L., 322
Chapin, J. W., 1229
Chareonviriyaphap, T., 545
Charlet, L. D., 1248, 1555
Chaudhury, M. F., 241
Chauhan, K. R., 1751
Chen, Q., 1450
Chen, W.-L., 685
Chen, W. M., 1854
Chen, Y., 1821
Cherry, S., 381
Chiasson, H., 459
Chicca, M., 1637
Chilcutt, C. F., 88, 327, 1428
Childers, C. C., 233
Choo, H. Y., 103
Chouinard, G., 737
Cinanni, L., 1047
Civolani, S., 1637
Clark, E. L., 1304
Clark, T. L., 557
Clarke, S. R., 1298
Clement, S. L., 131
Cloyd, R. A., 459, 1353

Coker, C. E. H., 1400
Colbo, M., 838
Coleman, P. G., 1642
Contreras, J., 916
Cook, D., 1916
Cook, S., 381
Coombs, J. J., 573
Cooper, S. G., 573
Cormier, D., 737
Corrales, C. E., 1781
Correa, J. C., 327
Coy, A. E., 1704
Crickmore, N., 932
Cross, J. V., 745
Crosslin, J. M., 656
Crowder, D. W., 1650
Cullen, E. M., 1258
Cutler, G. C., 765

D

Dalirsefat, S. B., 939
Dang, P. M., 1033
Danka, R. G., 267
Dashiell, K. E., 1464
Dastur, S. R., 1871
Daves, C. A., 969, 976
Davidson, S. A., 781
Davis, F. M., 969, 976
de Lame, F. M., 1316
Del Estal, P., 773, 1789
Dennehy, T. J., 1650
Deshmukh, S. D., 357
de Sousa-Majer, M. J., 1416
Diaz-Fleischer, F., 1153
Diaz-Montano, J., 984, 1657
Dickinson, A., 1353
Dietrich, G., 622
DiFonzo, C. D., 1258
Dindo, M. L., 26
Dionne, J., 1344
Dixon, A. F. G., 1781
Dixon, P. L., 838
Dolan, M. C., 622
Dong, P., 1476
Donnelly, C. A., 1642
Doolittle, M., 880
Dosdall, L. M., 215
Douches, D. S., 573
Doukas, D., 110, 389
Dowd, S. E., 1033
Dowell, F. E., 759, 1714
Downes, S., 1844
Duan, C.-X., 1450
Duchon, S., 545

E

Edu, J., 273, 1180
Eger, J. E., 233
Eitam, A., 670
Elberson, L. R., 131
Ellers-Kirk, C., 1650
Ellsworth, P. C., 1650
El-Sayed, A. M., 745
Epsky, N. D., 580
Ericsson, J. D., 31
Erler, F., 66
Espada, C., 580
Esperk, T., 627
Espino, L., 1282
Etheridge, J. L., 1400
Etxeberria, E., 1938
Evans, M. A., 131

F

Fadel, W., 1945
Fang, J. C., 20
Fathipour, Y., 350, 1098
Fedina, T. Y., 1924
Feng, X., 909
Fernández, E., 916
Ferracini, C., 1910
Ferro, D. N., 1871
Flanders, K. L., 790
Flinn, P. W., 591
Flores, S., 291
Folkins, M., 880
Follett, P. A., 251, 1091, 1499, 1519
Forsline, P. L., 1663
Frank, K., 136
Franzen, L. D., 1692
Fraser, I., 1304
Fraser, S., 1938
French, B. W., 1542
Frérot, B., 1797
Froeba, J. G., 117
Fuentes-Contreras, E., 551

G

Gadau, J., 1033
Gahan, L. J., 187
Galal, N., 1871
Galetto, L., 1504
Galvan, T. L., 1000
Gao, C. F., 1854
Garsia, K. A., 894
Gaussoin, R., 136
Gebre-Amlak, A., 1248, 1555
George, J., 431
Gerard, P. D., 488
Gerry, A. C., 1489
Ghimire, M. N., 1732
Girod, V., 1
Glogoza, P. A., 1258
Goettel, M. S., 31
Goltapeh, E. M., 1098
Gore, J., 1569, 1916
Goto, C., 1075
Gould, F., 180, 187
Grafius, E. J., 573
Grafton-Cardwell, E. E., 1435
Grávalos, C., 916
Gray, M. E., 1104
Greenberg, L., 148
Greene, J., 1916
Grewal, P., 1604
Grieshop, M. J., 591
Grillenberger, B., 1033
Grubbs, M. A., 619
Gu, H., 375
Guillén, M., 664
Guldin, J., 381
Guo, H. F., 20
Guo, Y., 909, 1409
Gupta, G., 830
Gut, L. J., 1316, 1360
Gutsche, A. R., 1692

H

Haas, M., 1360
Haghani, M., 350
Hall, D. G., 233, 847
Hall, D. R., 745
Hallett, R. H., 1335
Hansen, L. S., 1011
Hardie, D. C., 1416
Haverty, M. I., 1381
He, X., 49
He, Y. P., 1854
Heath, R. R., 580
Heckel, D. G., 187
Hellier, B. C., 131
Hellmich, R. L., 1129
Henderson, G., 866
Heng-Moss, T. M., 136, 1692
Henneberry, T. J., 1053
Hennessey, M. K., 251
Hentz, M. G., 1887
Heraty, J. M., 664
Herbert, D. A., Jr., 1241
Hernández-Crespo, P., 1836
Hesler, L. S., 1464, 1496
Hibbard, B. E., 557
Higdon, M. L., 557
Higgins, R. A., 1221
Higgins, T. J. V., 1416
Higley, L. G., 1268, 1692
Hill, J. H., 808
Hilton, R., 702
Hirose, E., 680
Hodgson, E. W., 1258
Hoffmann, A. A., 1741
Hogsette, J. A., 483
Hollaway, L., 1672
Hollingsworth, R. G., 1519
Hollomon, C. Z., 1400
Holst, N., 1
Hou, C., 202
Hou, M., 1160
Hoy, C. W., 526
Hu, F., 1476
Huang, F., 164, 1862, 1880
Huang, L. Q., 1854
Huang, Y., 202, 1672
Hubbard, J. L., 790
Hubert, J., 586, 1928

Hull, L. A., 421
Humes, K., 381
Hunt, T. E., 1258, 1268
Hunter, W. B., 1033
Hurst, T. P., 61
Husseneder, C., 1037, 1370
Hutchison, W. D., 1000
Hwang, F.-C., 1166
Hwang, S.-Y., 1166

I

Il'ichev, A. L., 1360
Isaacs, R., 855, 1596
Izadi, H., 361

J

Jackai, L. E. N., 315
Jackson, D. M., 566
Jackson, R. E., 180, 1569
Jallow, M. F.A., 526
James, S. S., 307
Jang, E. B., 1124
Jeffery, J. A. L., 1512
Jenkins, P. E., 855
Jervis, M., 1215
Jia, F., 759
Jia, F.-X., 1476
Jiang, X.-F., 1525
Jirakanjanakit, N., 545
Johnson, D. T., 398
Johnson, J. A., 1482
Johnson, K. D., 1258
Johnson, P. D., 1241
Johnson, T. A., 1328
Jonas, V. M., 1821
Jordan, D. L., 1241
Juba, T. R., 801
Judson, A. K., 1033
Jung, S.-C., 246

K

Kabaluk, J. T., 31
Kabashima, J. N., 148
Kaewpa, D., 946
Kalender, Y., 1533
Kamali, K., 1098
Karchesy, J. J., 622
Kavallieratos, N. G., 599, 1724
Kay, B. H., 61, 1512
Kaya, H. K., 103
Keddie, B. A., 215
Kehrli, P., 11
Keller, J. J., 258
Kemp, K. E., 1353
Kemper, C. J., 155
Kendra, P. E., 580
Kheradmand, K., 1098
Khosla, R., 781
Kim, Y.-G., 246
Klein, M. G., 307
Knight, A. L., 702
Knight, P. R., 1400
Knodel, J. J., 710
Kocourek, F., 1587
Kohama, T., 730
Konstantopoulou, M. A., 1215
Kopittke, R. A., 1136
Krakowsky, M. D., 1704
Krawczyk, G., 421
Krischik, V. A., 830, 1611
Krishnareddy, K. B., 903
Krizkova-Kudlikova, I., 1928
Kumano, N., 730
Kumar, P., 411
Kumar, R., 619
Kuruvinashetti, M. S., 903

L

Lacasa, A., 123
Lacey, L. A., 702
Lafleur, G., 737
Lam, W.-K. F., 823
Lambkin, T. A., 1136, 1423
Landolt, P. J., 366
Langemeier, M. R., 604
Lanzoni, A., 298
Lapchin, L., 225
Lapitan, N. L. V., 990
Lapointe, S. L., 847
Larentzaki, E., 1194
Láscares, S., 1173
Laurent, P., 1797
Laurin, M.-C., 1210
Lax, A., 880
Leblanc, L., 670
Lee, C.-Y., 467
Lee, D. W., 103
Lee, R. D., 1704
Lee, S. M., 103
Lee, V. S., 467
Lefebvre, D., 1756
Lei, C.-L., 1450
Leis, M., 1637
Leonard, B. R., 164, 1560, 1880
Leoncini, P., 1504
Leopold, R. A., 685
Leskey, T. C., 343, 1663
Levi, V., 1751
Lewis, B. A., 398
Lewis, L. C., 1129
Lewis, S. M., 1924
Li, H., 1862
Li, M., 202
Li, Y.-C., 990
Liang, G., 909
Liburd, O. E., 1622
Liedo, P., 291
Liu, B. S., 20
Liu, J.-L., 375
Liu, S. Q., 1298
Liu, X. G., 1854
Liu, Y.-B., 717
Lloyd, J. E., 619
Lo, P., 745
Lockwood, M. E., 1400
Löfstedt, C., 1017
López, J. D., Jr., 187
Lord, J. C., 1071
Lorenz, G., 1916
Louis, J., 984
Lowery, D. T., 1831
Lu, W., 1160
Luo, L.-Z., 1525
Luo, S., 909
Luppichini, P., 1391
Luttrell, R. G., 921

M

Ma, E., 1409
Ma, P.W.K., 976
Mackey, B. E., 723
MacRae, I. V., 1258
Maehara, N., 49
Magaña, C., 1836
Maghirang, E., 759, 1714
Mahon, R. J., 894, 1844
Maier, C. T., 1291
Malo, E. A., 1173
Malone, S., 1241
Malvar, R. A., 209
Manachini, B., 1047
Mangan, R. L., 440
Mankin, R. W., 790
Manning, L.-A., 745
Mannion, C. M., 307, 405
Mao, L., 866
Marchal, C., 225
Marchetti, E., 26
Marcus, M. A., 180
Margolies, D. C., 1821, 1904
Martin, P. A. W., 680
Martin, R. A., 283
Martínez, A.-M., 773
Marzachì, C., 1504
Masetti, A., 298
Mastro, V., 1304
Matocha, J. E., 88
Mazomenos, B. E., 1215
McCornack, B. P., 1258
McCoy, C. M., 155
McCullough, D. G., 1304, 1577
McGhee, P., 1360
McInnis, D. O., 273, 723
McKern, J. A., 398
McMullen, M. D., 1470
McPherson, R. M., 962, 1456
Meagher, R. L., 954, 1569
Medina, P., 1789
Meikle, W. G., 1
Mercadier, G., 1
Meyer, J. A., 155
Meyer, R. F., 1248, 1555
Miao, X., 202
Micinski, S., 187
Mihok, S., 613
Miles, J. W., 1896

Miller, D. R., 815
Miller, F., 1808
Miller, J. R., 1316
Milonas, P. G., 195
Mirhoseini, S. Z., 939
Mitcham, E. J., 1084
Mohan, S., 976
Moharil, M. P., 357
Monroe, W. A., 976
Montenieri, J. A., 622
Montoya, P., 291
Moore, S. H., 1880
Morales, J. J., 1789
Moreno, D. S., 440
Mornhinweg, D. W., 1685
Morse, J. G., 1053, 1435
Mpakou, F. D., 1724
Mpassoukou, A. E., 1724
Mukawa, S., 1075
Mullens, B. A., 1489
Mulrooney, J. E., 488
Munyaneza, J. E., 656
Münzbergová, Z., 586
Musser, F. R., 1194, 1916
Myers, C. T., 421, 1663
Myers, J. H., 31

N

Nagoshi, R. N., 954, 1569
Nalepa, C. A., 1633
Nansen, C., 1
Naranjo, S. E., 1053
Nault, B. A., 1194
Navone, P., 298
Nay, J. E., 1773
Nechols, J. R., 591, 1821, 1904
Neven, L., 1499
Nguyen, S. N., 534
Ni, X., 1704
Niehuis, O., 1033
Nigg, H. N., 1938
Nimbalkar, S. A., 357
Nordlund, D. A., 283
Notz, A., 801
Nuessly, G. S., 1887
Nylin, S., 627

O

Odvody, G. N., 327
Ohno, S., 730
Okutani-Akamatsu, Y., 1276
Oliver, J. B., 307
Oliver, J. E., 1124
Olsen, K. M., 894, 1844
Olson, D. L., 710, 1201
Olson, J. K., 1282
O'Neal, M. E., 1258
O'Neil, R. J., 1258
Opit, G. P., 1821
Oppert, B., 1862
Ordás, A., 209
Ordás, B., 209
Ortego, F., 1836
Othman, A. S., 467
Oyediran, I. O., 557

P

Pabón, A., 1896
Pahio, E., 273, 1180
Paine, T. D., 148
Panella, N. A., 622
Park, C. G., 103
Park, J. K., 103
Park, Y.-L., 1773
Parsons, C. K., 838
Pasqualini, E., 1637
Patil, B. V., 903
Patrick, C., 1679
Payne, C. C., 110, 389
Peairs, F. B., 781
Pell, C., 1304
Peng, J., 990
Peretto, R., 1637
Perez-Mendoza, J., 1714
Perring, T. M., 1764, 1773
Peteinatos, G. G., 599
Peterson, C. J., 488
Petrou, S. E., 599
Phillips, T. W., 1732
Pierce, C.M.F., 1104
Pierre, J., 1756
Pineda, S., 773
Plate, J., 1194
Plummer, P. R., 155
Poehling, H.-M., 411
Poland, T. M., 1304
Poncet, C., 225
Porter, A. H., 1871
Porter, D. R., 1672, 1679, 1685
Potter, B. D., 1258
Potter, D. A., 431, 451
Pourkheirandish, M., 939
Power, K., 1604
Prabhaker, N., 1053
Puterka, G. J., 1679
Putoa, R., 670

Q

Quinto, V., 916

R

Raccah, B., 1504
Raghubeer, E., 1499
Ragsdale, D. W., 1258
Rahman, K., 1229
Raina, A., 880
Ramaswamy, S., 759
Rao, N.G.V., 357
Reagan, T. E., 54
Reardon, B. J., 1116, 1129
Reardon, K. T., 1129
Reay-Jones, F.P.F., 54
Reding, M. E., 307
Redmond, C. T., 431
Reese, J. C., 984, 1657
Reinke, M. D., 72
Remmers, J., 327
Revilla, P., 209
Reyes, M., 551
Ricci, M., 1781
Rice, M. E., 808
Rice, S. J., 1136, 1423
Riedell, W. E., 1542
Riegel, C., 1370
Rill, S., 1435
Ripa, R., 1391
Robacker, D. C., 1147
Robbins, J., 1916
Roberts, P. M., 1456
Robertson, J., 1945
Robinson, W. A., 509
Rodd, C., 273
Rodríguez, E., 172
Rojas, J. C., 1173
Rongnoparut, P., 545, 946
Royals, B. M., 1241
Royalty, R. N., 431
Rull, J., 752, 1153
Rust, M. K., 148, 495, 1391
Ryan, P. A., 61, 1512
Ryne, C., 1017

S

Sabater-Muñoz, B., 1444
Sadof, C. S., 95
Saengtharatip, S., 545
Samih, M. A., 361
San Andrés, V., 1444
Sanchez, J. A., 123
Sánchez-Ramos, I., 195
Sandanayaka, W. R. M., 745
Sappington, T. W., 1116
Saracco, P., 1504
Saran, R. K., 495
Sarath, G., 1692
Sarfraz, M., 215
Satpute, N. S., 357
Sauphanor, B., 551
Savopoulou-Soultani, M., 195
Sayyed, A. H., 932
Schapaugh, W. T., 984, 1657
Scharf, M. E., 534
Schneider, M.-I., 773
Schöller, M., 591
Schumann, R. A., 1938
Scott-Dupree, C. D., 765
Scully, B. T., 1887
Shaw, P. W., 745
Shelly, T. E., 273, 1180
Shelton, A. M., 1194
Shelton, T. G., 488
Shen, J. L., 1854
Shen, T.-C., 1166
Shi, G. L., 1298
Shimazu, M., 49
Shin, H. K., 103

Shufran, K., 1679
Shufran, K. A., 1685
Siegert, N. W., 1577
Silverman, J., 626
Silvie, P., 954
Simard, L., 1344
Simms, D. M., 1370
Simpson, T., 1084
Sims, S. R., 871
Siskos, E. P., 1215
Skoda, S. R., 241
Smagghe, G., 773
Smirle, M. J., 1831
Smith, D. B., 283
Smith, H. A., 39
Smith, R. A., 283
Smith, S., 1304
Smitley, D. R., 103
Snook, M. E., 1887
Snyder, D. E., 155
Soengas, P., 209
Song, C., 534
Sorenson, C. E., 801
Sotelo, G., 1896
Southern, P. S., 801
Spear-O'Mara, J., 335
Srinivasan, R., 646
Stanley, M. C., 509
Stará, J., 1587
Stark, J. D., 1027
Starkey, R., 155
Staska, R. T., 131
Stejskal, V., 586, 1928
Stelinski, L. L., 1360
Stewart, S., 1916
Stodola, T. J., 195
Stuart, R. J., 1938
Studebaker, G., 1916
Su, N., 1450
Su, N.-Y., 475, 1391
Subrahmanyam, B., 361
Suckling, D. M., 745
Sumerford, D. V., 1129
Summers, J., 258
Sumrall, B. H., 283
Sun, J. H., 1298
Sun, J.-Z., 517, 1400
Svensson, G. P., 1017
Swier, S. R., 1604
Symmes, E. J., 1764

T

Tabashnik, B. E., 1650
Talebi, A. A., 350
Tameru, B., 315
Tammaru, T., 627
Tarpy, D. R., 258
Tauer, C. G., 1672
Temple, J. H., 1560
Tenczar, E. G., 1611
Thomas, D. B., 1147
Thomas, J. S., 1229
Thomson, L. J., 1741
Throne, J. E., 1714
Tian, Y., 909
Tikar, S. N., 357
Tilley, D. R., 604
Tipping, C., 1084
Toews, M. D., 1714
Toledo, J., 291, 1173
Tomanović, Ž., 599
Tomic-Carruthers, N., 1062
Toscano, N. C., 1053
Trimble, R. M., 1815
Turner, N. C., 1416

U

Uchida, G. K., 723
Ueckermann, E. A., 1098
Upton, J. E., 656
Urbaneja, A., 664, 1444

V

Valderrama, A. M., 172
Valle, J., 773
van Achterberg, K., 298
Vandervoort, C., 1596
Van Duyn, J. W., 180
Van Mele, P., 695
van Tellingen, E., 695
Vargas, R. I., 322, 670, 723, 1027
Vayssières, J.-F., 695
Vázquez, A., 580
Velásquez, N., 172
Venette, R. C., 1258
Vijaykumar, 903
Villaseñor, A., 291
Vincent, C., 737
Viñuela, E., 773, 1789
Vitullo, J., 405
Vitullo, J. M., 95
Vrolijks, J., 695

W

Wagner, M. R., 1808
Wagner, T. L., 488
Walker, J. T. S., 745
Wallis, R., 745
Wan, J.-M., 1450
Wang, F., 375, 1188
Wang, J.-J., 1476
Wang, J.-K., 1450
Wang, J. P., 20
Wang, S., 405
Wang, Y. N., 1298
Watanabe, T., 1276
Way, M. O., 54, 1282
Wen, J., 1160
Wenninger, E. J., 847
Werren, J. H., 1033
Wheeler, M., 1945
White, W. H., 155
Widstrom, N. W., 1887
Wilde, G., 1221
Williams, C., 381
Williams, K. A., 1821
Williams, W. P., 969, 976, 1470
Williamson, R. C., 1328
Wilson, L. T., 54
Winkle, J. R., 155
Wise, J. C., 1596
Woolfolk, S. W., 283
Wratten, S. D., 11
Wu, H., 1409
Wu, J.-C., 375, 1188
Wu, K., 909
Wu, X., 1880
Wu, Y. Q., 1672

X

Xie, Z., 1409
Xu, A., 202
Xu, W., 1704
Xue, S., 375, 1188

Y

Yafuso, C. M., 1650
Yang, G.-Q., 1188
Yang, J. J., 1938
Yang, M., 1409
Yeap, B.-K., 467
Yegen, O., 66
Yoksan, S., 545
Young, L., 136
Young, S. R., 894
Youssef, N. N., 307
Yu, T. Q., 1298
Yu, Y.-S., 375, 1188

Z

Zack, R. S., 366
Zaidi, M. A., 172
Zalucki, M. P., 1136
Zapata, A., 172
Zeller, W., 66
Zhai, H.-Q., 1450
Zhang, A., 343, 405
Zhang, J., 909
Zhang, L., 1525
Zhang, Q.-H., 1751
Zhao, L. L., 1298
Zhong, W. F., 20
Zhou, W. J., 1854
Zhu, K. Y., 1862
Zhu, Y. C., 1854
Zurowski, C. L., 1831

CUMULATIVE KEY WORD INDEX
Journal of Economic Entomology
Volume 100, 2007

α

α-amylase inhibitor,
Bruchus pisorum, 1416
head capsule widths, 1416
Pisum sativum, 1416
transgenic peas, 1416

A

abamectin,
azadirachtin, 411
Bemisia tabaci, 411
persistence, 411
spinosad, 411
abamectin susceptibility,
Cacopsylla pyri, 1637
insecticide resistance, 1637
pest control, 1637
Acalymma vittatum,
Cucumis melo, 823
Diabrotica undecimpunctata howardi, 823
direct scouting, 823
yellow sticky trap, 823
acaricidal activity,
Juglans regia, 1298
Tetranychus cinnabarinus, 1298
Tetranychus viennensis, 1298
accuracy,
bias, 1916
efficiency, 1916
Lygus lineolaris, 1916
Acer rubrum,
chlorpyrifos, 307
nursery production, 307
plant immersion, 307
Popillia japonica, 307
acetamiprid,
inheritance, 932
Plutella xylostella, 932
pyrethroids, 932
resistance, 932
acetylcholinesterase, 1409, 1836
citrus, 1836
esterases, 1836
general esterase, 1409
glutathione S-transferase, 1409
insecticides, 1836
malathion, 1409
Oxya chinensis, 1409
synergists, 1836
Acrosternum hilare,
Euschistus servus, 1456
host plant resistance, 1456
Nezara viridula, 1456
Aculops pelekassi,
Phyllocoptruta oleivora, 233
presence–absence sampling, 233
proportions, 233
adult beetle,
microbial control, 49
nonwoven fabric, 49
pine wilt disease, 49
advisory,
integrated pest management, 1241
tobacco thrips, 1241
tomato spotted wilt virus, 1241
virginia market-type peanut, 1241
Aedes (*Stegomyia*),
diagnostic dose, 545
insecticide, 545
Thailand, 545
Aedes aegypti,
biocidal, 622
essential oils, 622
Ixodes scapularis, 622
Xenopsylla cheopis, 622
AFF lure,
Anastrepha ludens, 1147
attractants, 1147
BioLure, 1147
traps, 1147
AFLP,
gene flow, 1525
genetic diversity, 1525
geographic population, 1525
Mythimna separata, 1525
Africa,
CGIAR, 695
conservation biological control, 695
fruit flies, 695
mango, 695
aggregation behavior,
management, 1633
multicolored Asian lady beetle, 1633
negative phototaxis, 1633
urban pest, 1633
aggregation pheromone, 1751, 1924
aphids, 1751
biocontrol, 1751
dimethyldecanal, 1924
goldeneyed lacewing, 1751
stored products, 1924
trap response, 1924
Agrilus planipennis,
Fraxinus, 1304
invasive species, 1304
phytosanitary treatment, 1304
Agromyzidae,
conservation biological control, 298
ecological infrastructures, 298
hymenopteran parasitoids, 298
weeds, 298
AkseBio2,
BioZell2000-B, 66
botanical insecticide, 66
Cacopsylla pyri, 66
management, 66
alate monitoring,
Coptotermes formosanus, 1400
invasive species, 1400
swarming behaviors, 1400
termite survey, 1400
alates,
Formosan subterranean termite, 517
integrated pest management, 517
landscape mulch management, 517
allele frequency,
Bt resistance, 1844
Cry1Ac, 1844
F_2 screen, 1844
Helicoverpa armigera, 1844
allergy,
botanical insecticides, 586
food, 586
grain, 586
mite, 586
Alphitobius diaperinus,
darkling beetle, 1136
insect pest, 1136
poultry litter, 1136
Tenebrionidae, 1136
ammonia quantification,
Anastrepha ludens, 580
Ceratitis capitata, 580
Fourier transform spectroscopy, 580
Tephritidae, 580
amplified fragment length polymorphism markers,
Bombyx mori, 939
genetic diversity, 939
silkworm, 939

Anastrepha ludens, 291, 440, 580, 752, 1147, 1153
AFF lure, 1147
ammonia quantification, 580
attractants, 1147
Beauveria bassiana, 291
BioLure, 1147
Ceratitis capitata, 580
Fourier transform spectroscopy, 580
GF-120, 440
horizontal transmission, 291
irradiation doses, 1153
mating competitiveness, 752, 1153
Mexican fruit fly, 440
pathogenicity, 291
Solbait, 440
sterile insect technique, 752, 1153
sterility induction, 1153
strain replacement, 752
Tephritidae, 580
traps, 1147
virulence, 291
Anastrepha serpentina,
calling behavior, 1173
mass rearing, 1173
sterile insect technique, 1173
Anobiidae,
Bostrichidae, 1732
Bruchidae, 1732
Curculionidae, 1732
Gelichiidae, 1732
Anopheles minimus,
CYP6AA3, 946
deltamethrin, 946
kinetics, 946
NADPH-cytochrome P450 reductase, 946
Anoplura,
little blue cattle louse, 619
Solenopotes capillatus, 619
ant management,
bait preference, 509
crazy ants, 509
Paratrechina longicornis, 509
antagonism,
Bemisia tabaci, 886
organophosphates, 886
potentiation, 886
pyrethroids, 886
antibiosis, 215, 1464
antixenosis, 215, 1464
crucifers, 215
diamondback moth, 215
host plant resistance, 1464
PI 436684, 1464
PI 548664, 1464
tolerance, 215
Anticarsia gemmatalis, 962, 1229
groundnut, 1229
host plant resistance, 962
Hypena scabra, 962
insecticide efficacy, 1229
peanut, 1229
Pseudoplusia includens, 962
threecornered alfalfa hopper, 1229
tomato spotted wilt virus, 1229
antifeedant,
azadirachtin, 117
feeding deterrent, 117
grasshopper, 117
locust, 117
antioxidant enzymes,
Galleria mellonella, 1533
midgut, 1533
penicillin, 1533
transaminases, 1533
antixenosis, 215, 1464
antibiosis, 215, 1464
crucifers, 215
diamondback moth, 215
host plant resistance, 1464
PI 436684, 1464
PI 548664, 1464
tolerance, 215
Anystis baccarum,
carbamates, 1210
neonicotinoids, 1210
organophosphates, 1210
pyrethroids, 1210
Aonidiella aurantii,
baseline susceptibility, 1435
development, 1435
fecundity, 1435
insect growth regulator, 1435
aphid, 123, 1672, 1751
aggregation pheromone, 1751
biocontrol, 1751
density thresholds, 123
goldeneyed lacewing, 1751
host resistance, 1672
pepper, 123
quantitative trait loci, 1672
Schizaphis graminum, 1672
simple sequence repeats, 1672
tolerance, 123
aphid predators,
conservation biocontrol, 39
insectary crops, 39
organic agriculture, 39
Syrphidae, 39
Aphis glycines, 984, 1657
chlorophyll losses, 1657
feeding behavior, 984
Glycine max, 984, 1657
Apis mellifera, 1, 267
Beauveria bassiana, 1
foraging behavior, 267
hive treatment, 1
pollination, 267
Vaccinium angustifolium, 267
Varroa destructor, 1
apple, 421, 702, 737
Conotrachelus nenuphar, 737
CpGV, 702
developmental rate, 421
ethyl (*E,Z*)-2,4-decadienoate, 702
highbush blueberry, 737
kairomone, 702
oriental fruit moth, 421
peach, 421
pear, 702
population dynamics, 421
traps, 737
apple orchard,
codling moth, 1587
cross-resistance, 1587
Cydia pomonella, 1587
resistance, 1587
armyworm,
cytochrome oxidase I, 954
Spodoptera frugiperda, 954
aromatherapy,
Mediterranean fruit fly, 273
sterile insect technique, 273
artificial diet,
Dipel, 1221
Helicoverpa zea, 1221
larval behavior, 1221
ash utilization,
eradication, 1577
Fraxinus, 1577
invasive pest, 1577
nonindigenous species, 1577
Asiatic garden beetle,
Bacillus thuringiensis serovar *japonensis*, 1604
European chafer, 1604
Japanese beetle, 1604
oriental beetle, 1604
attract and kill,
attractant, 366
behavior, 366
feeding, 366
moth, 366
attractant, 366, 483, 1124, 1147
AFF lure, 1147
Anastrepha ludens, 1147
attract and kill, 366
Bactrocera, 1124
bait, 483
behavior, 366
BioLure, 1147
feeding, 366
house fly, 483
melon fly, 1124
moth, 366
raspberry ketone formate, 1124
semiochemicals, 1124

toxicant, 483
trap, 483, 1147
attraction,
barriers, 1353
fungus gnats, 1353
greenhouse, 1353
pest management, 1353
augmentative biological control,
habitat complexity, 591
retail stores, 591
stored-products, 591
azadirachtin, 95, 117, 411, 451
abamectin, 411
antifeedant, 117
Bemisia tabaci, 411
feeding deterrent, 117
feeding-induced aggregation, 95
grasshopper, 117
imidacloprid, 95
kaolin, 451
limonene, 451
locust, 117
persistence, 411
Popillia japonica, 95, 451
pyrethroid, 451
spinosad, 411
thresholds, 95
azinphos-methyl,
Cydia pomonella, 551
glutathione S-transferases, 551
insecticide, 551
resistance, 551

B

Bacillus thuringiensis, 172, 180, 195, 246, 573, 894, 921, 1129, 1642, 1862
Bt, 1129
Bt resistance, 894
cross-resistance, 180
Cry1Ac, 894
Cry2Ab, 894
Cry2Ab2, 921
cry1Ia1, 573
cry3A, 573
European corn borer, 1129
F_2 screen resistance management, 195
genetic transformation, 172
glandular trichomes, 573
Helicoverpa armigera, 894
Helicoverpa zea, 921
Heliothis virescens, 180, 921
high-dose/refuge strategy, 195, 1642
insect resistance, 172
LC_{50}, 921
mathematical modeling, 1642
monitoring, 1129
Ostrinia nubilalis, 1862
pathogenicity, 246
Plutella xylostella, 246
proteinase-mediated resistance, 1862
resistance management, 180, 1129, 1642, 1862
resistance mechanisms, 1862
RIDL, 1642
S. chacoense-derived resistance, 573
septicemia, 246
Sesamia nonagrioides, 195
Solanum tuberosum, 172
Tecia solanivora, 172
transgenic maize, 195
Vip3A, 180
Xenorhabdus nematophila, 246
Bacillus thuringiensis serovar *japonensis*,
Asiatic garden beetle, 1604
European chafer, 1604
Japanese beetle, 1604
oriental beetle, 1604
Bactericera cockerelli,
phytoplasma, 656
potato, 656
zebra chip, 656
Bactrocera,
attractants, 1124
melon fly, 1124
raspberry ketone formate, 1124
semiochemicals, 1124
Bactrocera dorsalis,
fruit fly rearing, 322
liquid diet, 322
wheat germ oil, 322
Bactrocera species,
classical biocontrol, 670
competitive displacement, 670
oriental fruit fly, 670
bait,
attractant, 483
house fly, 483
toxicant, 483
trap, 483
bait design,
bait station, 1938
Caribbean fruit fly, 1938
bait preference,
ant management, 509
crazy ants, 509
Paratrechina longicornis, 509
bait station,
bait design, 1938
Caribbean fruit fly, 1938
barley,
host plant resistance, 1679
host race, 1679
wheat, 1679
barriers,
attraction, 1353
fungus gnats, 1353
greenhouse, 1353
pest management, 1353
baseline susceptibility,
Aonidiella aurantii, 1435
development, 1435
fecundity, 1435
insect growth regulator, 1435
Beauveria bassiana, 1, 291, 1071
Anastrepha ludens, 291
Apis mellifera, 1
desiccation, 1071
flour beetle, 1071
hive treatment, 1
horizontal transmission, 291
humidity, 1071
pathogenicity, 291
Tribolium castaneum, 1071
Varroa destructor, 1
virulence, 291
bedding plants,
damage threshold, 1904
esthetic damage, 1904
floriculture, 1904
spider mites, 1904
behavior,
attract and kill, 366
attractant, 366
feeding, 366
moth, 366
behavioral avoidance,
permethrin, 526
Plutella xylostella, 526
pyrethroids, 526
behavioral resistance,
(Z)-9-tricosene, 1489
fly bait, 1489
house flies, 1489
methomyl, 1489
Bemisia tabaci, 411, 886, 1053, 1160, 1650
abamectin, 411
antagonism, 886
azadirachtin, 411
cucumber, 1160
genetics, 1650
haplodiploid, 1650
Homalodisca lacerta, 1053
Homalodisca vitripennis, 1053
insecticide impacts, 1053
insecticide resistance, 1650
organophosphates, 886
parasitoids, 1053
persistence, 411
potentiation, 886
pyrethroids, 886
pyriproxyfen, 1650
sampling, 1160
spinosad, 411
within-plant distribution, 1160
beneficial arthropods,
Blissus occiduus, 136
buffalograss, 136

sampling, 136
western chinch bugs, 136
beneficials,
IOBC, 1741
pesticide metrics, 1741
sustainable agriculture, 1741
bias,
accuracy, 1916
efficiency, 1916
Lygus lineolaris, 1916
binding,
cross-resistance, 909
Cry1Ac, 909
Cry2Ab, 909
Helicoverpa armigera, 909
bioassay,
Diabrotica, 566
Ipomoea, 566
plant resistance, 566
biocidal,
Aedes aegypti, 622
essential oils, 622
Ixodes scapularis, 622
Xenopsylla cheopis, 622
biocontrol, 680, 1027, 1751
aggregation pheromone, 1751
aphids, 1751
Diabrotica, 680
goldeneyed lacewing, 1751
IOBC, 1027
Nezara viridula, 680
pesticides, 1027
biological control, 11, 31, 283, 664
Cirrospilus coachellae, 664
citrus, 664
enzyme biomarkers, 11
insect rearing, 283
interaction, 31
lacewing, 283
life history, 664
Marmara gulosa, 664
Metarhizium, 31
MOD system, 283
mortality rates, 11
naturalyte, 31
pesticides, 11
Verticel larval rearing unit, 283
wireworm, 31
BioLure,
AFF lure, 1147
Anastrepha ludens, 1147
attractants, 1147
traps, 1147
biorational insecticides,
hybrid tea rose, 830
imidacloprid, 830
Popillia japonica, 830
biosecurity,
irradiation, 1519
Opogona sacchari, 1519
quarantine treatment, 1519
sterilizing dose, 1519
biotype, 990, 1685
Diuraphis noxia, 1685
Duraphis noxia, 990
elicitor, 990
Hordeum vulgare, 1685
host plant resistance, 1685
plant–aphid interaction, 990
Triticum aestivum, 990, 1685
BioZell2000-B,
AkseBio2, 66
botanical insecticide, 66
Cacopsylla pyri, 66
management, 66
blackberries,
entomopathogenic nematodes, 398
Heterorhabditis, 398
insecticides, 398
Steinernema, 398
blacklight trap,
mesh cone trap, 1797
monitoring, 1797
sticky trap, 1797
water pan trap, 1797
Blattella germanica,
surfactant, 871
synergism, 871
Blissus occiduus,
beneficial arthropods, 136
buffalograss, 136
sampling, 136
western chinch bugs, 136
blueberry, 1596, 1622
distribution, 1622
hot-spots, 1622
imidacloprid, 1596
Japanese beetle, 1596
PIC-Triad, 1596
sampling, 1622
sublethal, 1596
thrips, 1622
Bombus terrestris,
colony growth, 1756
drift, 1756
pollination, 1756
Bombyx mori, 202, 939
amplified fragment length polymorphism markers, 939
DNA fingerprinting, 202
genetic diversity, 939
intersimple sequence repeats, 202
silkworm, 939
Bostrichidae,
Anobiidae, 1732
Bruchidae, 1732
Curculionidae, 1732
Gelichiidae, 1732
botanic termiticide,
Ku Shen, 866
wood protectants, 866
botanical insecticide, 66, 586, 1215
AkseBio2, 66
allergy, 586
BioZell2000-B, 66
Cacopsylla pyri, 66
food, 586
grain, 586
management, 66
mite, 586
olive fly, 1215
plant extracts, 1215
Rutaceae, 1215
Brachiaria,
competition, 1896
multiple resistance, 1896
spittlebug, 1896
Brassicaceae,
Contarinia nasturtii, 1335
cruciferous vegetables, 1335
cruciferous weeds, 1335
host susceptibility, 1335
bread wheat,
Cephus cinctus, 79
economic loss, 79
sawfly, 79
yield loss, 79
breeding system, 1037, 1370
colony elimination, 1370
colony identification, 1037
microsatellite genotyping, 1037, 1370
movement, 1037
Rhinotermitidae, 1037
broiler house,
darkling beetle, 1423
insecticide resistance, 1423
lesser mealworm, 1423
Bruchidae,
Anobiidae, 1732
Bostrichidae, 1732
Curculionidae, 1732
Gelichiidae, 1732
Bruchus pisorum,
head capsule widths, 1416
Pisum sativum, 1416
transgenic peas, 1416
α-amylase inhibitor, 1416
Bt, 1116, 1129
Bacillus thuringiensis, 1129
dispersal, 1116
European corn borer, 1116, 1129
monitoring, 1129
Ostrinia nubilalis, 1116
resistance management, 1129
transgenic corn, 1116

Bt cotton,
crop strains, 903
crossability, 903
fecundity, 903
Helicoverpa armigera, 903
Bt maize,
Diatraea saccharalis, 164
insecticide resistance management, 164
resistance monitoring, 164
Bt resistance, 894, 1844
allele frequency, 1844
Bacillus thuringiensis, 894
Cry1Ac, 894, 1844
Cry2Ab, 894
F_2 screen, 1844
Helicoverpa armigera, 894, 1844
Bt resistance management,
Cry1Ab protein, 1880
Diatraea saccharalis, 1880
transgenic corn, 1880
buffalograss,
beneficial arthropods, 136
Blissus occiduus, 136
sampling, 136
western chinch bugs, 136

C

C. formosanus,
hexaflumuron, 475
R. flavipes, 475
Sentricon, 475
cabbage maggot,
diversification, 838
intercropping, 838
pest management, 838
relay cropping, 838
Cacopsylla pyri, 66, 1637
abamectin susceptibility, 1637
AkseBio2, 66
BioZell2000-B, 66
botanical insecticide, 66
insecticide resistance, 1637
management, 66
pest control, 1637
calling behavior,
Anastrepha serpentina, 1173
mass rearing, 1173
sterile insect technique, 1173
Calliphora vicina,
germplasm regeneration, 131
insect pollination, 131
leek, 131
Musca domestica, 131
canola, 710, 765, 1201
clothianidin, 765
flea beetle, 710, 1201
honey bees, 765
injury, 710
insecticides, 710, 1201
lethal time, 1201
long-term assessment, 765
relative potencies, 1201
seed-treatment, 765
yield, 710
carbamates,
Anystis baccarum, 1210
neonicotinoids, 1210
organophosphates, 1210
pyrethroids, 1210
carbon dioxide,
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
Pseudococcus maritimus, 1084
quarantine, 1084
Caribbean fruit fly,
bait design, 1938
bait station, 1938
cellulose fiber,
larval diet, 241
rearing, 241
screwworm, 241
spray-dried blood, 241
Cephus cinctus,
bread wheat, 79
economic loss, 79
sawfly, 79
yield loss, 79
Ceratitis capitata, 580, 1180, 1444
ammonia quantification, 580
Anastrepha ludens, 580
Fourier transform spectroscopy, 580
male age, 1180
mating success, 1180, 1444
Mediterranean fruit fly, 1180
polymerase chain reaction-restriction fragment length polymorphism, 1444
sperm, 1444
spermathecae, 1444
sterile insect technique, 1180
Tephritidae, 580
cereal pest,
geographical distribution, 1781
host preference, 1781
Sipha maydis, 1781
CGIAR,
Africa, 695
conservation biological control, 695
fruit flies, 695
mango, 695
Chilo suppressalis, 375, 1188, 1854
fipronil, 1854
imidacloprid, 375, 1188
juvenile hormone, 1188
juvenile hormone III, 375
molting hormone, 375, 1188
monosultap, 1854
resistance, 1854
rice plant, 375
triazophos, 1854
chipping,
Saperda vestita, 1328
systemic insecticides, 1328
Tilia spp., 1328
chlorophyll losses,
Aphis glycines, 1657
Glycine max, 1657
chlorpyrifos,
Acer rubrum, 307
nursery production, 307
plant immersion, 307
Popillia japonica, 307
Chrysanthemum yellows phytoplasma,
Euscelidius variegatus, 1504
Euscelis incisus, 1504
Macrosteles quadripunctulatus, 1504
vector transmission, 1504
Chrysomela scripta,
hybrid poplar, 1611
imidacloprid, 1611
novel application methods, 1611
Cirrospilus coachellae,
biological control, 664
citrus, 664
life history, 664
Marmara gulosa, 664
citrus, 664, 1836
acetylcholinesterase, 1836
biological control, 664
Cirrospilus coachellae, 664
esterases, 1836
insecticides, 1836
life history, 664
Marmara gulosa, 664
synergists, 1836
classical biocontrol,
Bactrocera species, 670
competitive displacement, 670
oriental fruit fly, 670
Cleridae,
Conophthorus coniperda, 815
Enoclerus nigripes, 815
kairomone, 815
Scolytidae, 815
clone,
mustard, 1764
okra, 1764
transmission efficiency, 1764
clothianidin,
canola, 765
honey bees, 765
long-term assessment, 765
seed-treatment, 765
codling moth, 1499, 1587
apple orchard, 1587
cross-resistance, 1587
Cydia pomonella, 1499, 1587

quarantine treatment, 1499
resistance, 1587
Rhagoletis indifferens, 1499
western cherry fruit fly, 1499
COII mitochondrial DNA,
Coptotermes gestroi, 467
Coptotermes vastator, 467
ribosomal DNA, 467
synonymity, 467
cold storage,
egg parasitoid, 685
fitness, 685
glassy-winged sharpshooter, 685
host suitability, 685
cold treatment,
dried fruits, 1482
Indianmeal moth, 1482
navel orangeworm, 1482
tree nuts, 1482
colony density,
colony size, 1381
cuticular hydrocarbons, 1381
mark–release–recapture, 1381
soldier proportions, 1381
colony elimination,
breeding system, 1370
microsatellite genotyping, 1370
colony growth,
Bombus terrestris, 1756
drift, 1756
pollination, 1756
colony identification,
breeding system, 1037
microsatellite genotyping, 1037
movement, 1037
Rhinotermitidae, 1037
colony size,
colony density, 1381
cuticular hydrocarbons, 1381
mark–release–recapture, 1381
soldier proportions, 1381
community ecology,
Cry1Ab, 327
genetically modified, 327
maize, 327
plant resistance, 327
competition, 148, 1896
Brachiaria, 1896
interspecific aggression, 148
Linepithema humile, 148
multiple resistance, 1896
Solenopsis invicta, 148
spittlebug, 1896
competitive displacement,
Bactrocera species, 670
classical biocontrol, 670
oriental fruit fly, 670
Conophthorus coniperda,
Cleridae, 815
Enoclerus nigripes, 815
kairomone, 815
Scolytidae, 815
Conotrachelus nenuphar,
apple, 737
highbush blueberry, 737
traps, 737
conservation biocontrol,
aphid predators, 39
insectary crops, 39
organic agriculture, 39
Syrphidae, 39
conservation biological control, 298, 695
Africa, 695
Agromyzidae, 298
CGIAR, 695
ecological infrastructures, 298
fruit flies, 695
hymenopteran parasitoids, 298
mango, 695
weeds, 298
conservation tillage,
Helicoverpa zea, 88
Oebalus pugnax, 88
rice stink bug, 88
sorghum headworm, 88
contaminant pest,
Harmonia axyridis, 1000
resampling software, 1000
wine grapes, 1000
Contarinia nasturtii,
Brassicaceae, 1335
cruciferous vegetables, 1335
cruciferous weeds, 1335
host susceptibility, 1335
continuous in vitro culture,
hatching, 26
macrotype eggs, 26
oviposition substrate, 26
controlled atmosphere, 717, 1084
carbon dioxide, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
postharvest quality, 717
Pseudococcus maritimus, 1084
quarantine, 1084
quarantine treatment, 717
ultralow oxygen, 717
western flower thrips, 717
Coptotermes formosanus, 880, 1400
alate monitoring, 1400
d-limonene, 880
fumigation, 880
invasive species, 1400
orange oil extract, 880
swarming behaviors, 1400
termite survey, 1400
tunneling behavior, 880
Coptotermes gestroi,
COII mitochondrial DNA, 467
Coptotermes vastator, 467
ribosomal DNA, 467
synonymity, 467
Coptotermes vastator,
COII mitochondrial DNA, 467
Coptotermes gestroi, 467
ribosomal DNA, 467
synonymity, 467
corn,
management zones, 781
site-specific management, 781
cotton, 187, 1560, 1569
DNA diagnostics, 187
fall armyworm, 1569
heat units, 1560
host strains, 1569
injury levels, 1560
integrated pest management, 1560
resistance monitoring, 187
southern green stink bug, 1560
Spodoptera frugiperda, 1569
stable isotope, 1569
tobacco budworm, 187
cotton aphid,
cotton bollworm, 1428
cotton fleahopper, 1428
sweetpotato whitefly, 1428
western flower thrips, 1428
cotton bollworm,
cotton aphid, 1428
cotton fleahopper, 1428
sweetpotato whitefly, 1428
western flower thrips, 1428
cotton fleahopper,
cotton aphid, 1428
cotton bollworm, 1428
sweetpotato whitefly, 1428
western flower thrips, 1428
counting,
Cucumis sativus, 225
Frankliniella occidentalis, 225
projection pursuit regression, 225
Rosa × *hybrida*, 225
coverage probabilities,
delta method, 1945
probit regression, 1945
CpGV,
apple, 702
ethyl (*E*,*Z*)-2,4-decadienoate, 702
kairomone, 702
pear, 702
crazy ants,
ant management, 509
bait preference, 509
Paratrechina longicornis, 509

crop strains,
Bt cotton, 903
crossability, 903
fecundity, 903
Helicoverpa armigera, 903
cross-resistance, 180, 909, 1587
apple orchard, 1587
Bacillus thuringiensis, 180
binding, 909
codling moth, 1587
Cry1Ac, 909
Cry2Ab, 909
Cydia pomonella, 1587
Helicoverpa armigera, 909
Heliothis virescens, 180
resistance, 1587
resistance management, 180
Vip3A, 180
crossability,
Bt cotton, 903
crop strains, 903
fecundity, 903
Helicoverpa armigera, 903
cruciferous vegetables,
Brassicaceae, 1335
Contarinia nasturtii, 1335
cruciferous weeds, 1335
host susceptibility, 1335
cruciferous weeds,
Brassicaceae, 1335
Contarinia nasturtii, 1335
cruciferous vegetables, 1335
host susceptibility, 1335
crucifers,
antibiosis, 215
antixenosis, 215
diamondback moth, 215
tolerance, 215
Cry1Ab,
community ecology, 327
genetically modified, 327
maize, 327
plant resistance, 327
Cry1Ab protein,
Bt resistance management, 1880
Diatraea saccharalis, 1880
transgenic corn, 1880
Cry1Ac, 894, 909, 1844
allele frequency, 1844
Bacillus thuringiensis, 894
binding, 909
Bt resistance, 894, 1844
cross-resistance, 909
Cry2Ab, 894, 909
F_2 screen, 1844
Helicoverpa armigera, 894, 909, 1844
cry1Ia1,
Bacillus thuringiensis (Bt), 573
cry3A, 573
glandular trichomes, 573
S. chacoense-derived resistance, 573
Cry2Ab, 894, 909
Bacillus thuringiensis, 894
binding, 909
Bt resistance, 894
cross-resistance, 909
Cry1Ac, 894, 909
Helicoverpa armigera, 894, 909
Cry2Ab2,
Bacillus thuringiensis, 921
Helicoverpa zea, 921
Heliothis virescens, 921
LC_{50}, 921
cry3A,
Bacillus thuringiensis (Bt), 573
cry1Ia1, 573
glandular trichomes, 573
S. chacoense-derived resistance, 573
Cry3Bb1,
Diabrotica virgifera virgifera, 557
glyphosate, 557
grassy weeds, 557
herbicide, 557
cucumber, 350, 1160
Bemisia tabaci, 1160
degree-days, 350
development, 350
Liriomyza sativae, 350
sampling, 1160
temperature dependent, 350
within-plant distribution, 1160
Cucumis melo,
Acalymma vittatum, 823
Diabrotica undecimpunctata howardi, 823
direct scouting, 823
yellow sticky trap, 823
Cucumis sativus,
counting, 225
Frankliniella occidentalis, 225
projection pursuit regression, 225
Rosa × *hybrida*, 225
cultural management,
Ectomyelois ceratoniae, 1773
geographic information system, 1773
SADIE, 1773
Curculionidae,
Anobiidae, 1732
Bostrichidae, 1732
Bruchidae, 1732
Gelichiidae, 1732
cuticular hydrocarbons,
colony density, 1381
colony size, 1381
mark–release–recapture, 1381
soldier proportions, 1381
Cydia pomonella, 551, 1499, 1587
apple orchard, 1587
azinphos-methyl, 551
codling moth, 1499, 1587
cross-resistance, 1587
glutathione *S*-transferases, 551
insecticide, 551
quarantine treatment, 1499
resistance, 551, 1587
Rhagoletis indifferens, 1499
western cherry fruit fly, 1499
Cylas formicarius elegantulus, 315, 730
dispersal ability, 730
gamma radiation, 730
quantitative risk analysis, 315
sterile insect technique, 730
sweet potato, 315
sweetpotato weevil, 730
Cylindrocopturus adspersus, 1248, 1555
Dectes texanus, 1248, 1555
Pelochrista womonana, 1248, 1555
pest management, 1248, 1555
sunflower, 1248, 1555
Cyp12D1,
Cyp6g1, 534
cytochrome P450, 534
fumigant, 534
negative cross-resistance, 534
CYP6AA3,
Anopheles minimus, 946
deltamethrin, 946
kinetics, 946
NADPH-cytochrome P450 reductase, 946
Cyp6g1,
Cyp12D1, 534
cytochrome P450, 534
fumigant, 534
negative cross-resistance, 534
cytochrome oxidase I,
armyworm, 954
Spodoptera frugiperda, 954
cytochrome P450,
Cyp12D1, 534
Cyp6g1, 534
fumigant, 534
negative cross-resistance, 534

D

d-limonene,
Coptotermes formosanus, 880
fumigation, 880
orange oil extract, 880
tunneling behavior, 880
damage relationships,
flue-cured tobacco, 801
infestation, 801
treatment threshold, 801

damage threshold,
bedding plants, 1904
esthetic damage, 1904
floriculture, 1904
spider mites, 1904
darkling beetle, 1136, 1423
Alphitobius diaperinus, 1136
broiler house, 1423
insect pest, 1136
insecticide resistance, 1423
lesser mealworm, 1423
poultry litter, 1136
Tenebrionidae, 1136
Dectes texanus, 1248, 1555
Cylindrocopturus adspersus, 1248, 1555
Pelochrista womonana, 1248, 1555
pest management, 1248, 1555
sunflower, 1248, 1555
defoliation,
Pyrrhalta luteola, 1808
Ulmus spp., 1808
degree-days, 335, 350, 1098
cucumber, 350
development, 350
Heterocampa guttivitta, 335
Liriomyza sativae, 350
nonlinear models, 1098
pheromone trapping, 335
population monitoring, 335
Sancassania rodionovi, 1098
sugar maple, 335
temperature dependent, 350
temperature-dependent development, 1098
delayed toxicity,
nonrepellency, 495
Reticulitermes hesperus, 495
termite movement, 495
delta method,
coverage probabilities, 1945
probit regression, 1945
deltamethrin,
Anopheles minimus, 946
CYP6AA3, 946
kinetics, 946
NADPH-cytochrome P450 reductase, 946
density thresholds,
aphids, 123
pepper, 123
tolerance, 123
desiccation,
Beauveria bassiana, 1071
flour beetle, 1071
humidity, 1071
Tribolium castaneum, 1071
development, 350, 1435, 1476
Aonidiella aurantii, 1435
baseline susceptibility, 1435
cucumber, 350
degree-days, 350
fecundity, 1435
insect growth regulator, 1435
life table parameters, 1476
Liposcelis tricolor, 1476
Liriomyza sativae, 350
reproduction, 1476
temperature dependent, 350
Wolbachia, 1476
developmental diet breadth,
feeding performance, 1166
host plant range, 1166
Lymantria xylina, 1166
developmental polymorphism,
instar number, 627
life history, 627
plasticity, 627
threshold size, 627
developmental rate,
apple, 421
oriental fruit moth, 421
peach, 421
population dynamics, 421
Diabrotica, 566, 680, 1542
bioassay, 566
biocontrol, 680
IPM, 1542
Ipomoea, 566
landscape, 1542
Nezara viridula, 680
plant resistance, 566
semiochemical baits, 1542
Diabrotica undecimpunctata howardi,
Acalymma vittatum, 823
Cucumis melo, 823
direct scouting, 823
yellow sticky trap, 823
Diabrotica virgifera virgifera,
Cry3Bb1, 557
glyphosate, 557
grassy weeds, 557
herbicide, 557
diagnostic dose,
Aedes (*Stegomyia*), 545
insecticide, 545
Thailand, 545
diamondback moth,
antibiosis, 215
antixenosis, 215
crucifers, 215
tolerance, 215
diatomaceous earth, 599, 1724
laboratory bioassays, 1724
mortality, 599
progeny production, 599
Prostephanus truncatus, 599
Rhyzopertha dominica, 1724
Sitophilus oryzae, 1724
stored maize, 599
Tribolium confusum, 1724
Diatraea saccharalis, 164, 1880
Bt maize, 164
Bt resistance management, 1880
Cry1Ab protein, 1880
insecticide resistance management, 164
resistance monitoring, 164
transgenic corn, 1880
diet,
nutrition, 1062
rearing, 1062
root, 1062
weevil, 1062
diffusion,
dispersal, 110
Encarsia formosa, 110
UV-blocking films, 110
dimethyldecanal,
aggregation pheromone, 1924
stored products, 1924
trap response, 1924
Dipel,
artificial diet, 1221
Helicoverpa zea, 1221
larval behavior, 1221
direct scouting,
Acalymma vittatum, 823
Cucumis melo, 823
Diabrotica undecimpunctata howardi, 823
yellow sticky trap, 823
dispersal, 110, 745, 1116
Bt, 1116
diffusion, 110
Encarsia formosa, 110
European corn borer, 1116
mass trapping, 745
multiple mating, 745
Ostrinia nubilalis, 1116
pheromone, 745
suppression, 745
transgenic corn, 1116
UV-blocking films, 110
dispersal ability,
Cylas formicarius elegantulus, 730
gamma radiation, 730
sterile insect technique, 730
sweetpotato weevil, 730
distribution,
blueberries, 1622
hot-spots, 1622
sampling, 1622
thrips, 1622
Diuraphis noxia,
biotype, 1685
Hordeum vulgare, 1685
host plant resistance, 1685
Triticum aestivum, 1685

diversification,
cabbage maggot, 838
intercropping, 838
pest management, 838
relay cropping, 838
DNA diagnostics,
cotton, 187
resistance monitoring, 187
tobacco budworm, 187
DNA fingerprinting,
Bombyx mori, 202
intersimple sequence repeats, 202
dominance,
insecticide resistance management, 1871
Leptinotarsa decemlineata, 1871
life history, 1871
dried fruits,
cold treatment, 1482
Indianmeal moth, 1482
navel orangeworm, 1482
tree nuts, 1482
drift,
Bombus terrestris, 1756
colony growth, 1756
pollination, 1756
Duraphis noxia,
biotype, 990
elicitor, 990
plant–aphid interaction, 990
Triticum aestivum, 990

E

Earias vitella,
insecticide toxicity, 357
temperature, 357
ecological infrastructures,
Agromyzidae, 298
conservation biological control, 298
hymenopteran parasitoids, 298
weeds, 298
economic loss,
bread wheat, 79
Cephus cinctus, 79
sawfly, 79
yield loss, 79
Ecotilling,
genome mapping, 1033
Nasonia, 1033
species identification, 1033
Ectomyelois ceratoniae,
cultural management, 1773
geographic information system, 1773
SADIE, 1773
efficiency,
accuracy, 1916
bias, 1916
Lygus lineolaris, 1916
egg parasitoid,
cold storage, 685
fitness, 685
glassy-winged sharpshooter, 685
host suitability, 685
elicitor,
biotype, 990
Duraphis noxia, 990
plant–aphid interaction, 990
Triticum aestivum, 990
ELISA,
extraction, 1928
grain, 1928
mite, 1928
storage, 1928
emergence, 1104, 1194
maize, 1104
onions, 1194
overwintering, 1194
phenology, 1104
soybean, 1104
Thrips tabaci, 1194
volunteer onions, 1194
emergence inhibition,
insecticides, 1512
sublethal, 1512
survival, 1512
emulsified wax,
Grapholita molesta, 1316
mating disruption, 1316
pheromone dispenser, 1316
Encarsia formosa,
diffusion, 110
dispersal, 110
UV-blocking films, 110
Endopiza viteana,
insect growth regulator, 855
methoxyfenozide, 855
Paralobesia viteana, 855
spinosad, 855
Enoclerus nigripes,
Cleridae, 815
Conophthorus coniperda, 815
kairomone, 815
Scolytidae, 815
entomopathogenic nematodes,
blackberries, 398
Heterorhabditis, 398
insecticides, 398
Steinernema, 398
enzyme biomarkers,
biological control, 11
mortality rates, 11
pesticides, 11
enzyme-linked immunosorbent assay,
Fabaceae, 808
insect herbivory, 808
leaf area measurement, 808
reverse transcription-polymerase chain reaction, 808
Eoreuma loftini,
Mexican rice borer, 54
movement, 54
pheromone trap, 54
eradication,
ash utilization, 1577
Fraxinus, 1577
invasive pest, 1577
nonindigenous species, 1577
essential oils, 459, 622
Aedes aegypti, 622
biocidal, 622
fungus gnats, 459
Ixodes scapularis, 622
mealybugs, 459
pest management, 459
western flower thrips, 459
Xenopsylla cheopis, 622
esterases,
acetylcholinesterase, 1836
citrus, 1836
insecticides, 1836
synergists, 1836
esthetic damage,
bedding plants, 1904
damage threshold, 1904
floriculture, 1904
spider mites, 1904
ethyl (*E,Z*)-2,4-decadienoate,
apple, 702
CpGV, 702
kairomone, 702
pear, 702
ethylene glycol,
mineral oil, 723
nontarget insects, 723
preservatives, 723
propylene glycol, 723
European chafer,
Asiatic garden beetle, 1604
Bacillus thuringiensis serovar *japonensis*, 1604
Japanese beetle, 1604
oriental beetle, 1604
European corn borer, 1116, 1129
Bacillus thuringiensis, 1129
Bt, 1116, 1129
dispersal, 1116
monitoring, 1129
Ostrinia nubilalis, 1116
resistance management, 1129
transgenic corn, 1116
Euscelidius variegatus,
Chrysanthemum yellows phytoplasma, 1504
Euscelis incisus, 1504
Macrosteles quadripunctulatus, 1504
vector transmission, 1504

Euscelis incisus,
Chrysanthemum yellows phytoplasma, 1504
Euscelidius variegatus, 1504
Macrosteles quadripunctulatus, 1504
vector transmission, 1504
Euschistus servus,
Acrosternum hilare, 1456
host plant resistance, 1456
Nezara viridula, 1456
exotic beetle,
host preference, 1291
Juniperus, 1291
Thuja, 1291
trap log, 1291
extraction,
ELISA, 1928
grain, 1928
mite, 1928
storage, 1928

F

F_2 screen,
allele frequency, 1844
Bt resistance, 1844
Cry1Ac, 1844
Helicoverpa armigera, 1844
F_2 screen resistance management,
Bacillus thuringiensis, 195
high-dose/refuge strategy, 195
Sesamia nonagrioides, 195
transgenic maize, 195
Fabaceae,
enzyme-linked immunosorbent assay, 808
insect herbivory, 808
leaf area measurement, 808
reverse transcription-polymerase chain reaction, 808
fall armyworm, 1470, 1569, 1887
cotton, 1569
host strains, 1569
insect resistance, 1470
maize, 1470
maysin, 1887
Primetime, 1887
QTL, 1470
southwestern corn borer, 1470
Spodoptera frugiperda, 1569
stable isotope, 1569
Zapalote Chico, 1887
fecundity, 646, 903, 1435
Aonidiella aurantii, 1435
baseline susceptibility, 1435
Bt cotton, 903
crop strains, 903
crossability, 903
development, 1435
Helicoverpa armigera, 903
host preference, 646
insect growth regulator, 1435
mixed viral infections, 646
potato, 646
feeding,
attract and kill, 366
attractant, 366
behavior, 366
moth, 366
feeding behavior,
Aphis glycines, 984
Glycine max, 984
feeding deterrent,
antifeedant, 117
azadirachtin, 117
grasshopper, 117
locust, 117
feeding lure,
Korea, 103
pheromone traps, 103
Popillia japonica, 103
Popillia quadriguttata, 103
turfgrass, 103
feeding performance,
developmental diet breadth, 1166
host plant range, 1166
Lymantria xylina, 1166
feeding-induced aggregation,
azadirachtin, 95
imidacloprid, 95
Popillia japonica, 95
thresholds, 95
female attractiveness,
ovarian development, 1276
sex pheromone, 1276
Stenotus rubrovittatus, 1276
field sampling,
leafminer, 1910
pest management, 1910
Taylor's power law, 1910
urban green, 1910
field studies,
mating disruption, 1017
portable EAG, 1017
Pyralidae, 1017
stored-product pests, 1017
fipronil,
Chilo suppressalis, 1854
monosultap, 1854
resistance, 1854
triazophos, 1854
fitness,
cold storage, 685
egg parasitoid, 685
glassy-winged sharpshooter, 685
host suitability, 685
flea beetle, 710, 1201
canola, 710, 1201
injury, 710
insecticides, 710, 1201
lethal time, 1201
relative potencies, 1201
yield, 710
floriculture,
bedding plants, 1904
damage threshold, 1904
esthetic damage, 1904
spider mites, 1904
flour beetle,
Beauveria bassiana, 1071
desiccation, 1071
humidity, 1071
Tribolium castaneum, 1071
flue-cured tobacco,
damage relationships, 801
infestation, 801
treatment threshold, 801
fluorescent brightener,
granulovirus, 1075
Mamestra brassicae, 1075
nucleopolyhedrovirus, 1075
viral enhancement, 1075
fly bait,
(Z)-9-tricosene, 1489
behavioral resistance, 1489
house flies, 1489
methomyl, 1489
food,
allergy, 586
botanical insecticides, 586
grain, 586
mite, 586
foraging behavior,
Apis mellifera, 267
pollination, 267
Vaccinium angustifolium, 267
Formosan subterranean termite,
alates, 517
integrated pest management, 517
landscape mulch management, 517
Fourier transform spectroscopy,
ammonia quantification, 580
Anastrepha ludens, 580
Ceratitis capitata, 580
Tephritidae, 580
Frankliniella occidentalis, 225, 1084
carbon dioxide, 1084
controlled atmosphere, 1084
counting, 225
Cucumis sativus, 225
Platynota stultana, 1084
projection pursuit regression, 225
Pseudococcus maritimus, 1084
quarantine, 1084
Rosa × *hybrida*, 225

Fraxinus, 1304, 1577
Agrilus planipennis, 1304
ash utilization, 1577
eradication, 1577
invasive pest, 1577
invasive species, 1304
nonindigenous species, 1577
phytosanitary treatment, 1304
fruit flies, 251, 695, 1091
Africa, 695
CGIAR, 695
conservation biological control, 695
Litchi chinensis, 1091
mango, 695
phytosanitary treatments, 251
postharvest treatment, 1091
probit 9, 251
quarantine pests, 1091
quarantine statistics, 251
fruit fly rearing,
Bactrocera dorsalis, 322
liquid diet, 322
wheat germ oil, 322
fumigant,
Cyp12D1, 534
Cyp6g1, 534
cytochrome P450, 534
negative cross-resistance, 534
fumigation,
Coptotermes formosanus, 880
d-limonene, 880
orange oil extract, 880
tunneling behavior, 880
fungus gnats, 459, 1353
attraction, 1353
barriers, 1353
essential oils, 459
greenhouse, 1353
mealybugs, 459
pest management, 459, 1353
western flower thrips, 459

G

Galleria mellonella,
antioxidant enzymes, 1533
midgut, 1533
penicillin, 1533
transaminases, 1533
gamma radiation,
Cylas formicarius elegantulus, 730
dispersal ability, 730
sterile insect technique, 730
sweetpotato weevil, 730
gas exchange,
host plant resistance, 1692
photosynthesis, 1692
plant–insect interaction, 1692
Gelichiidae,
Anobiidae, 1732
Bostrichidae, 1732
Bruchidae, 1732
Curculionidae, 1732
gene flow,
AFLP, 1525
genetic diversity, 1525
geographic population, 1525
Mythimna separata, 1525
general esterase,
acetylcholinesterase, 1409
glutathione *S*-transferase, 1409
malathion, 1409
Oxya chinensis, 1409
genetic diversity, 939, 1525
AFLP, 1525
amplified fragment length polymorphism markers, 939
Bombyx mori, 939
gene flow, 1525
geographic population, 1525
Mythimna separata, 1525
silkworm, 939
genetic transformation,
Bacillus thuringiensis, 172
insect resistance, 172
Solanum tuberosum, 172
Tecia solanivora, 172
genetically modified,
community ecology, 327
Cry1Ab, 327
maize, 327
plant resistance, 327
genetics,
Bemisia tabaci, 1650
haplodiploid, 1650
insecticide resistance, 1650
pyriproxyfen, 1650
genome mapping,
Ecotilling, 1033
Nasonia, 1033
species identification, 1033
geographic information system, 381, 1773
cultural management, 1773
Ectomyelois ceratoniae, 1773
hazard rating, 381
Landsat ETM+, 381
SADIE, 1773
southern pine beetle, 381
geographic population,
AFLP, 1525
gene flow, 1525
genetic diversity, 1525
Mythimna separata, 1525
geographical distribution,
cereal pest, 1781
host preference, 1781
Sipha maydis, 1781
geospatial mapping,
geostatistics, 790
SADIE, 790
soil insects, 790
sound, 790
geostatistics,
geospatial mapping, 790
SADIE, 790
soil insects, 790
sound, 790
germplasm regeneration,
Calliphora vicina, 131
insect pollination, 131
leek, 131
Musca domestica, 131
GF-120,
Anastrepha ludens, 440
Mexican fruit fly, 440
Solbait, 440
glandular trichomes,
Bacillus thuringiensis (Bt), 573
cry1Ia1, 573
cry3A, 573
S. chacoense-derived resistance, 573
glassy-winged sharpshooter,
cold storage, 685
egg parasitoid, 685
fitness, 685
host suitability, 685
glutathione *S*-transferase,
acetylcholinesterase, 1409
general esterase, 1409
malathion, 1409
Oxya chinensis, 1409
glutathione *S*-transferases,
azinphos-methyl, 551
Cydia pomonella, 551
insecticide, 551
resistance, 551
Glycine max, 984, 1657
Aphis glycines, 984, 1657
chlorophyll losses, 1657
feeding behavior, 984
glyphosate,
Cry3Bb1, 557
Diabrotica virgifera virgifera, 557
grassy weeds, 557
herbicide, 557
goldeneyed lacewing,
aggregation pheromone, 1751
aphids, 1751
biocontrol, 1751
golf course,
Listronotus maculicollis, 1344
seasonal ecology, 1344
turfgrass insect, 1344
grain, 586, 1928
allergy, 586
botanical insecticides, 586
ELISA, 1928
extraction, 1928
food, 586
mite, 586, 1928
storage, 1928

granulovirus,
fluorescent brightener, 1075
Mamestra brassicae, 1075
nucleopolyhedrovirus, 1075
viral enhancement, 1075
grape berry moth,
mating disruption, 1815
pheromone, 1815
Grapholita molesta, 72, 1316
emulsified wax, 1316
mating disruption, 1316
methoxyfenozide, 72
pheromone dispenser, 1316
sexual behavior, 72
wind tunnel, 72
grasshopper,
antifeedant, 117
azadirachtin, 117
feeding deterrent, 117
locust, 117
grassy weeds,
Cry3Bb1, 557
Diabrotica virgifera virgifera, 557
glyphosate, 557
herbicide, 557
Greenhead trap,
Manitoba trap, 613
Nzi trap, 613
stable fly, 613
tabanids, 613
greenhouse,
attraction, 1353
barriers, 1353
fungus gnats, 1353
pest management, 1353
greening disease,
huanglongbing, 847
kaolin, 847
phenology, 847
Surround, 847
groundnut,
Anticarsia gemmatalis, 1229
insecticide efficacy, 1229
peanut, 1229
threecornered alfalfa hopper, 1229
tomato spotted wilt virus, 1229

H

habitat complexity,
augmentative biological control, 591
retail stores, 591
stored-products, 591
haplodiploid,
Bemisia tabaci, 1650
genetics, 1650
insecticide resistance, 1650
pyriproxyfen, 1650
Harmonia axyridis,
contaminant pest, 1000
resampling software, 1000
wine grapes, 1000
hatching,
continuous in vitro culture, 26
macrotype eggs, 26
oviposition substrate, 26
hazard rating,
geographic information system, 381
Landsat ETM+, 381
southern pine beetle, 381
head capsule widths,
Bruchus pisorum, 1416
Pisum sativum, 1416
transgenic peas, 1416
α-amylase inhibitor, 1416
heat treatment,
Rhyzopertha dominica, 604
risk analysis, 604
Sitophilus oryzae, 604
Tribolium castaneum, 604
heat units,
cotton, 1560
injury levels, 1560
integrated pest management, 1560
southern green stink bug, 1560
Helicoverpa armigera, 894, 903, 909, 1844
allele frequency, 1844
Bacillus thuringiensis, 894
binding, 909
Bt cotton, 903
Bt resistance, 894, 1844
crop strains, 903
cross-resistance, 909
crossability, 903
Cry1Ac, 894, 909, 1844
Cry2Ab, 894, 909
F_2 screen, 1844
fecundity, 903
Helicoverpa zea, 88, 759, 921, 1221, 1704
artificial diet, 1221
Bacillus thuringiensis, 921
conservation tillage, 88
Cry2Ab2, 921
Dipel, 1221
Heliothis virescens, 759, 921
husk extension, 1704
larval behavior, 1221
LC_{50}, 921
Oebalus pugnax, 88
rice stink bug, 88
silk-feeding bioassay, 1704
Sitophilus zeamais, 1704
sorghum headworm, 88
species identification, 759
stink bugs, 1704
Heliothis virescens, 180, 759, 921
Bacillus thuringiensis, 180, 921
cross-resistance, 180
Cry2Ab2, 921
Helicoverpa zea, 759, 921
LC_{50}, 921
resistance management, 180
species identification, 759
Vip3A, 180
herbaceous species,
holocycle, 1047
sharka, 1047
vector, 1047
weeds, 1047
herbicide,
Cry3Bb1, 557
Diabrotica virgifera virgifera, 557
glyphosate, 557
grassy weeds, 557
Heterocampa guttivitta,
degree-days, 335
pheromone trapping, 335
population monitoring, 335
sugar maple, 335
Heterorhabditis,
blackberries, 398
entomopathogenic nematodes, 398
insecticides, 398
Steinernema, 398
hexaflumuron,
C. formosanus, 475
R. flavipes, 475
Sentricon, 475
high-dose/refuge strategy, 195, 1642
Bacillus thuringiensis, 195, 1642
F_2 screen resistance management, 195
mathematical modeling, 1642
resistance management, 1642
RIDL, 1642
Sesamia nonagrioides, 195
transgenic maize, 195
highbush blueberry,
apple, 737
Conotrachelus nenuphar, 737
traps, 737
hive treatment,
Apis mellifera, 1
Beauveria bassiana, 1
Varroa destructor, 1
holocycle,
herbaceous species, 1047
sharka, 1047
vector, 1047
weeds, 1047
Homalodisca lacerta,
Bemisia tabaci, 1053
Homalodisca vitripennis, 1053
insecticide impacts, 1053

parasitoids, 1053
Homalodisca vitripennis,
Bemisia tabaci, 1053
Homalodisca lacerta, 1053
insecticide impacts, 1053
parasitoids, 1053
honey bees, 258, 765
canola, 765
clothianidin, 765
integrative pest management, 258
long-term assessment, 765
Russian stock. 258
seed-treatment, 765
tracheal mite, 258
varroa mite, 258
Hordeum vulgare,
biotype, 1685
Diuraphis noxia, 1685
host plant resistance, 1685
Triticum aestivum, 1685
horizontal transmission,
Anastrepha ludens, 291
Beauveria bassiana, 291
pathogenicity, 291
virulence, 291
host plant range,
developmental diet breadth, 1166
feeding performance, 1166
Lymantria xylina, 1166
host plant resistance, 962, 969, 976, 1456, 1464, 1679, 1685, 1692
Acrosternum hilare, 1456
antibiosis, 1464
Anticarsia gemmatalis, 962
antixenosis, 1464
barley, 1679
biotype, 1685
Diuraphis noxia, 1685
Euschistus servus, 1456
gas exchange, 1692
Hordeum vulgare, 1685
host race, 1679
Hypena scabra, 962
Nezara viridula, 1456
peritrophic membrane, 976
photosynthesis, 1692
PI 436684, 1464
PI 548664, 1464
plant–insect interaction, 1692
Pseudoplusia includens, 962
southwestern corn borer, 969, 976
Triticum aestivum, 1685
wheat, 1679
host preference, 646, 1291, 1781
cereal pest, 1781
exotic beetle, 1291
fecundity, 646
geographical distribution, 1781
Juniperus, 1291
mixed viral infections, 646
potato, 646
Sipha maydis, 1781
Thuja, 1291
trap log, 1291
host race,
barley, 1679
host plant resistance, 1679
wheat, 1679
host resistance,
aphid, 1672
quantitative trait loci, 1672
Schizaphis graminum, 1672
simple sequence repeats, 1672
host strains,
cotton, 1569
fall armyworm, 1569
Spodoptera frugiperda, 1569
stable isotope, 1569
host suitability,
cold storage, 685
egg parasitoid, 685
fitness, 685
glassy-winged sharpshooter, 685
host susceptibility,
Brassicaceae, 1335
Contarinia nasturtii, 1335
cruciferous vegetables, 1335
cruciferous weeds, 1335
hot-spots,
blueberries, 1622
distribution, 1622
sampling, 1622
thrips, 1622
house fly, 155, 483, 1489
attractant, 483
bait, 483
behavioral resistance, 1489
fly bait, 1489
imidacloprid, 155
methomyl, 155, 1489
Musca domestica, 155
spinosad, 155
toxicant, 483
trap, 483
(Z)-9-tricosene, 1489
huanglongbing,
greening disease, 847
kaolin, 847
phenology, 847
Surround, 847
humidity,
Beauveria bassiana, 1071
desiccation, 1071
flour beetle, 1071
Tribolium castaneum, 1071
husk extension,
Helicoverpa zea, 1704
silk-feeding bioassay, 1704
Sitophilus zeamais, 1704
stink bugs, 1704
hybrid poplar,
Chrysomela scripta, 1611
imidacloprid, 1611
novel application methods, 1611
hybrid tea rose,
biorational insecticides, 830
imidacloprid, 830
Popillia japonica, 830
hymenopteran parasitoids,
Agromyzidae, 298
conservation biological control, 298
ecological infrastructures, 298
weeds, 298
Hypena scabra,
Anticarsia gemmatalis, 962
host plant resistance, 962
Pseudoplusia includens, 962
Hyposoter didymator,
insecticides, 1789
protected stages, 1789

I

imidacloprid, 95, 155, 375, 830, 1188, 1596, 1611
azadirachtin, 95
biorational insecticides, 830
blueberry, 1596
Chilo suppressalis, 375, 1188
Chrysomela scripta, 1611
feeding-induced aggregation, 95
house fly, 155
hybrid poplar, 1611
hybrid tea rose, 830
Japanese beetle, 1596
juvenile hormone, 1188
juvenile hormone III, 375
methomyl, 155
molting hormone, 375, 1188
Musca domestica, 155
novel application methods, 1611
PIC-Triad, 1596
Popillia japonica, 95, 830
rice plant, 375
spinosad, 155
sublethal, 1596
thresholds, 95
Indianmeal moth,
cold treatment, 1482
dried fruits, 1482
navel orangeworm, 1482
tree nuts, 1482
infestation,
damage relationships, 801
flue-cured tobacco, 801
treatment threshold, 801

inheritance, 916, 932
acetamiprid, 932
insecticide resistance, 916
Plutella xylostella, 932
pyrethroids, 932
resistance, 932
Thysanoptera, 916
injury,
canola, 710
flea beetle, 710
insecticides, 710
yield, 710
injury levels,
cotton, 1560
heat units, 1560
integrated pest management, 1560
southern green stink bug, 1560
insect fragments,
internal insect infestation, 1714
near-infrared spectroscopy (NIRS), 1714
rice weevil, 1714
stored products, 1714
insect growth regulator, 361, 855, 1435
Aonidiella aurantii, 1435
baseline susceptibility, 1435
development, 1435
Endopiza viteana, 855
fecundity, 1435
methoxyfenozide, 855
ovicide, 361
oxime ethers, 361
Paralobesia viteana, 855
spinosad, 855
Spodoptera litura, 361
synergism, 361
insect herbivory,
enzyme-linked immunosorbent assay, 808
Fabaceae, 808
leaf area measurement, 808
reverse transcription-polymerase chain reaction, 808
insect pest,
Alphitobius diaperinus, 1136
darkling beetle, 1136
poultry litter, 1136
Tenebrionidae, 1136
insect pollination,
Calliphora vicina, 131
germplasm regeneration, 131
leek, 131
Musca domestica, 131
insect rearing,
biological control, 283
lacewing, 283
MOD system, 283
Verticel larval rearing unit, 283
insect resistance, 172, 1470
Bacillus thuringiensis, 172
fall armyworm, 1470
genetic transformation, 172
maize, 1470
QTL, 1470
Solanum tuberosum, 172
southwestern corn borer, 1470
Tecia solanivora, 172
insectary crops,
aphid predators, 39
conservation biocontrol, 39
organic agriculture, 39
Syrphidae, 39
insecticide, 398, 545, 551, 710, 1201, 1512, 1789, 1836
acetylcholinesterase, 1836
Aedes (*Stegomyia*), 545
azinphos-methyl, 551
blackberries, 398
canola, 710, 1201
citrus, 1836
Cydia pomonella, 551
diagnostic dose, 545
emergence inhibition, 1512
entomopathogenic nematodes, 398
esterases, 1836
flea beetle, 710, 1201
glutathione *S*-transferases, 551
Heterorhabditis, 398
Hyposoter didymator, 1789
injury, 710
lethal time, 1201
protected stages, 1789
relative potencies, 1201
resistance, 551
Steinernema, 398
sublethal, 1512
survival, 1512
synergists, 1836
Thailand, 545
yield, 710
insecticide efficacy,
Anticarsia gemmatalis, 1229
groundnut, 1229
peanut, 1229
threecornered alfalfa hopper, 1229
tomato spotted wilt virus, 1229
insecticide impacts,
Bemisia tabaci, 1053
Homalodisca lacerta, 1053
Homalodisca vitripennis, 1053
parasitoids, 1053
insecticide mixtures,
insecticide resistance, 1831
integrated pest management, 1831
kaolin, 1831
obliquebanded leafroller, 1831
insecticide resistance, 916, 1423, 1637, 1650, 1831
abamectin susceptibility, 1637
Bemisia tabaci, 1650
broiler house, 1423
Cacopsylla pyri, 1637
darkling beetle, 1423
genetics, 1650
haplodiploid, 1650
inheritance, 916
insecticide mixtures, 1831
integrated pest management, 1831
kaolin, 1831
lesser mealworm, 1423
obliquebanded leafroller, 1831
pest control, 1637
pyriproxyfen, 1650
Thysanoptera, 916
insecticide resistance management, 164, 1871
Bt maize, 164
Diatraea saccharalis, 164
dominance, 1871
Leptinotarsa decemlineata, 1871
life history, 1871
resistance monitoring, 164
insecticide toxicity,
Earias vitella, 357
temperature, 357
instar number,
developmental polymorphism, 627
life history, 627
plasticity, 627
threshold size, 627
integrated crop management,
IPM, 1821
Pelargonium peltatum, 1821
plant nutrition, 1821
Tetranychus urticae, 1821
integrated pest management, 517, 1241, 1560, 1831
advisory, 1241
alates, 517
cotton, 1560
Formosan subterranean termite, 517
heat units, 1560
injury levels, 1560
insecticide mixtures, 1831
insecticide resistance, 1831
kaolin, 1831
landscape mulch management, 517
obliquebanded leafroller, 1831
southern green stink bug, 1560
tobacco thrips, 1241
tomato spotted wilt virus, 1241

virginia market-type peanut, 1241
integrative pest management,
honey bees, 258
Russian stock, 258
tracheal mite, 258
varroa mite, 258
interaction,
biological control, 31
Metarhizium, 31
naturalyte, 31
wireworm, 31
intercropping,
cabbage maggot, 838
diversification, 838
pest management, 838
relay cropping, 838
internal insect infestation,
insect fragments, 1714
near-infrared spectroscopy (NIRS), 1714
rice weevil, 1714
stored products, 1714
intersimple sequence repeats,
Bombyx mori, 202
DNA fingerprinting, 202
interspecific aggression,
competition, 148
Linepithema humile, 148
Solenopsis invicta, 148
introduced pest,
natural enemies, 1268
Orius insidiosus, 1268
invasive pest,
ash utilization, 1577
eradication, 1577
Fraxinus, 1577
nonindigenous species, 1577
invasive species, 1258, 1304, 1400
Agrilus planipennis, 1304
alate monitoring, 1400
Coptotermes formosanus, 1400
Fraxinus, 1304
phytosanitary treatment, 1304
population dynamics, 1258
swarming behaviors, 1400
termite survey, 1400
yield loss, 1258
IOBC, 1027, 1741
beneficials, 1741
biocontrol, 1027
pesticide metrics, 1741
pesticides, 1027
sustainable agriculture, 1741
IPM, 1542, 1821
Diabrotica, 1542
integrated crop management, 1821
landscape, 1542
Pelargonium peltatum, 1821
plant nutrition, 1821
semiochemical baits, 1542
Tetranychus urticae, 1821
Ipomoea,
bioassay, 566
Diabrotica, 566
plant resistance, 566
irradiation,
biosecurity, 1519
Opogona sacchari, 1519
quarantine treatment, 1519
sterilizing dose, 1519
irradiation doses,
Anastrepha ludens, 1153
mating competitiveness, 1153
sterile insect technique, 1153
sterility induction, 1153
Ixodes scapularis,
Aedes aegypti, 622
biocidal, 622
essential oils, 622
Xenopsylla cheopis, 622

J

Japanese beetle, 1596, 1604
Asiatic garden beetle, 1604
Bacillus thuringiensis serovar *japonensis*, 1604
blueberry, 1596
European chafer, 1604
imidacloprid, 1596
oriental beetle, 1604
PIC-Triad, 1596
sublethal, 1596
Juglans regia,
acaricidal activity, 1298
Tetranychus cinnabarinus, 1298
Tetranychus viennensis, 1298
Juniperus,
exotic beetle, 1291
host preference, 1291
Thuja, 1291
trap log, 1291
juvenile hormone,
Chilo suppressalis, 1188
imidacloprid, 1188
molting hormone, 1188
juvenile hormone III,
Chilo suppressalis, 375
imidacloprid, 375
molting hormone, 375
rice plant, 375

K

kairomone, 702, 815
apple, 702
Cleridae, 815
Conophthorus coniperda, 815
CpGV, 702
Enoclerus nigripes, 815
ethyl (*E,Z*)-2,4-decadienoate, 702
pear, 702
Scolytidae, 815
kaolin, 451, 847, 1831
azadirachtin, 451
greening disease, 847
huanglongbing, 847
insecticide mixtures, 1831
insecticide resistance, 1831
integrated pest management, 1831
limonene, 451
obliquebanded leafroller, 1831
phenology, 847
Popillia japonica, 451
pyrethroid, 451
Surround, 847
kinetics,
Anopheles minimus, 946
CYP6AA3, 946
deltamethrin, 946
NADPH-cytochrome P450 reductase, 946
Korea,
feeding lure, 103
pheromone traps, 103
Popillia japonica, 103
Popillia quadriguttata, 103
turfgrass, 103
Ku Shen,
botanic termiticide, 866
wood protectants, 866

L

laboratory bioassays,
diatomaceous earths, 1724
Rhyzopertha dominica, 1724
Sitophilus oryzae, 1724
Tribolium confusum, 1724
lacewing,
biological control, 283
insect rearing, 283
MOD system, 283
Verticel larval rearing unit, 283
Landsat ETM+,
geographic information system, 381
hazard rating, 381
southern pine beetle, 381
landscape,
Diabrotica, 1542
IPM, 1542
semiochemical baits, 1542
landscape mulch management,
alates, 517
Formosan subterranean termite, 517
integrated pest management, 517
Laodelphax striatellus,
quantitative trait loci, 1450
recombinant inbred line population, 1450
resistance, 1450

rice, 1450
Lariophagus distinguendus,
life table, 1011
parasitoid-induced mortality, 1011
Sitophilus granarius, 1011
temperature, 1011
larval behavior,
artificial diet, 1221
Dipel, 1221
Helicoverpa zea, 1221
larval diet,
cellulose fiber, 241
rearing, 241
screwworm, 241
spray-dried blood, 241
larvicides,
Melanotaenia, 61
mosquito, 61
organophosphate, 61
sublethal effects, 61
LC_{50},
Bacillus thuringiensis, 921
Cry2Ab2, 921
Helicoverpa zea, 921
Heliothis virescens, 921
leaf area measurement,
enzyme-linked immunosorbent assay, 808
Fabaceae, 808
insect herbivory, 808
reverse transcription-polymerase chain reaction, 808
leafminer,
field sampling, 1910
pest management, 1910
Taylor's power law, 1910
urban green, 1910
leek,
Calliphora vicina, 131
germplasm regeneration, 131
insect pollination, 131
Musca domestica, 131
Leptinotarsa decemlineata,
dominance, 1871
insecticide resistance management, 1871
life history, 1871
lesser mealworm,
broiler house, 1423
darkling beetle, 1423
insecticide resistance, 1423
lethal time,
canola, 1201
flea beetle, 1201
insecticides, 1201
relative potencies, 1201
life history, 627, 664, 1871
biological control, 664
Cirrospilus coachellae, 664
citrus, 664
developmental polymorphism, 627
dominance, 1871
insecticide resistance management, 1871
instar number, 627
Leptinotarsa decemlineata, 1871
Marmara gulosa, 664
plasticity, 627
threshold size, 627
life table,
Lariophagus distinguendus, 1011
parasitoid-induced mortality, 1011
Sitophilus granarius, 1011
temperature, 1011
life table parameters,
development, 1476
Liposcelis tricolor, 1476
reproduction, 1476
Wolbachia, 1476
light-diffusion,
Trialeurodes vaporariorum, 389
UV-blocking films, 389
whitefly, 389
limonene,
azadirachtin, 451
kaolin, 451
Popillia japonica, 451
pyrethroid, 451
Linepithema humile,
competition, 148
interspecific aggression, 148
Solenopsis invicta, 148
Liposcelis tricolor,
development, 1476
life table parameters, 1476
reproduction, 1476
Wolbachia, 1476
liquid diet,
Bactrocera dorsalis, 322
fruit fly rearing, 322
wheat germ oil, 322
Liriomyza sativae,
cucumber, 350
degree-days, 350
development, 350
temperature dependent, 350
Listronotus maculicollis,
golf course, 1344
seasonal ecology, 1344
turfgrass insect, 1344
Litchi chinensis,
fruit flies, 1091
postharvest treatment, 1091
quarantine pests, 1091
little blue cattle louse,
Anoplura, 619
Solenopotes capillatus, 619
locust,
antifeedant, 117
azadirachtin, 117
feeding deterrent, 117
grasshopper, 117
long-term assessment,
canola, 765
clothianidin, 765
honey bees, 765
seed-treatment, 765
lures,
monitoring, 343
plum curculio, 343
release rate, 343
temperature, 343
Lygus lineolaris,
accuracy, 1916
bias, 1916
efficiency, 1916
Lymantria xylina,
developmental diet breadth, 1166
feeding performance, 1166
host plant range, 1166

M

Maconellicoccus hirsutus,
monitoring, 405
sex pheromone, 405
trapping, 405
Macrosteles quadripunctulatus,
Chrysanthemum yellows phytoplasma, 1504
Euscelidius variegatus, 1504
Euscelis incisus, 1504
vector transmission, 1504
macrotype eggs,
continuous in vitro culture, 26
hatching, 26
oviposition substrate, 26
maize, 327, 1104, 1470
community ecology, 327
Cry1Ab, 327
emergence, 1104
fall armyworm, 1470
genetically modified, 327
insect resistance, 1470
phenology, 1104
plant resistance, 327
QTL, 1470
southwestern corn borer, 1470
soybean, 1104
malathion,
acetylcholinesterase, 1409
general esterase, 1409
glutathione *S*-transferase, 1409
Oxya chinensis, 1409
male age,
Ceratitis capitata, 1180
mating success, 1180
Mediterranean fruit fly, 1180
sterile insect technique, 1180

Mamestra brassicae,
fluorescent brightener, 1075
granulovirus, 1075
nucleopolyhedrovirus, 1075
viral enhancement, 1075
management, 66, 1633
aggregation behavior, 1633
AkseBio2, 66
BioZell2000-B, 66
botanical insecticide, 66
Cacopsylla pyri, 66
multicolored Asian lady beetle, 1633
negative phototaxis, 1633
urban pest, 1633
management zones,
corn, 781
site-specific management, 781
mango,
Africa, 695
CGIAR, 695
conservation biological control, 695
fruit flies, 695
Manitoba trap,
Greenhead trap, 613
Nzi trap, 613
stable fly, 613
tabanids, 613
mark–release–recapture,
colony density, 1381
colony size, 1381
cuticular hydrocarbons, 1381
soldier proportions, 1381
Marmara gulosa,
biological control, 664
Cirrospilus coachellae, 664
citrus, 664
life history, 664
mass rearing,
Anastrepha serpentina, 1173
calling behavior, 1173
sterile insect technique, 1173
mass trapping,
dispersal, 745
multiple mating, 745
pheromone, 745
suppression, 745
mathematical modeling,
Bacillus thuringiensis, 1642
high-dose/refuge strategy, 1642
resistance management, 1642
RIDL, 1642
mating competitiveness, 752, 1153
Anastrepha ludens, 752, 1153
irradiation doses, 1153
sterile insect technique, 752, 1153
sterility induction, 1153
strain replacement, 752
mating disruption, 1017, 1316, 1360, 1815
emulsified wax, 1316
field studies, 1017
grape berry moth, 1815
Grapholita molesta, 1316
pheromone, 1815
pheromone dispenser, 1316
portable EAG, 1017
Pyralidae, 1017
sprayable microencapsulated pheromone, 1360
stored-product pests, 1017
Tortricidae, 1360
mating success, 1180, 1444
Ceratitis capitata, 1180, 1444
male age, 1180
Mediterranean fruit fly, 1180
polymerase chain reaction-restriction fragment length polymorphism, 1444
sperm, 1444
spermathecae, 1444
sterile insect technique, 1180
maysin,
fall armyworm, 1887
Primetime, 1887
Zapalote Chico, 1887
mealybugs,
essential oils, 459
fungus gnats, 459
pest management, 459
western flower thrips, 459
Mediterranean fruit fly, 273, 1180
aromatherapy, 273
Ceratitis capitata, 1180
male age, 1180
mating success, 1180
sterile insect technique, 273, 1180
Melanotaenia,
larvicides, 61
mosquito, 61
organophosphate, 61
sublethal effects, 61
melon fly,
attractants, 1124
Bactrocera, 1124
raspberry ketone formate, 1124
semiochemicals, 1124
mesh cone trap,
blacklight trap, 1797
monitoring, 1797
sticky trap, 1797
water pan trap, 1797
Metarhizium,
biological control, 31
interaction, 31
naturalyte, 31
wireworm, 31
methomyl, 155, 1489
(Z)-9-tricosene, 1489
behavioral resistance, 1489
fly bait, 1489
house flies, 1489
house fly, 155
imidacloprid, 155
Musca domestica, 155
spinosad, 155
methoxyfenozide, 72, 773, 855
Endopiza viteana, 855
Grapholita molesta, 72
insect growth regulator, 855
Paralobesia viteana, 855
pesticide residues, 773
reproductive parameters, 773
sexual behavior, 72
spinosad, 773, 855
Spodoptera littoralis, 773
wind tunnel, 72
Mexican fruit fly,
Anastrepha ludens, 440
GF-120, 440
Solbait, 440
Mexican rice borer,
Eoreuma loftini, 54
movement, 54
pheromone trap, 54
microbial control,
adult beetle, 49
nonwoven fabric, 49
pine wilt disease, 49
microsatellite genotyping, 1037, 1370
breeding system, 1037, 1370
colony elimination, 1370
colony identification, 1037
movement, 1037
Rhinotermitidae, 1037
midgut,
antioxidant enzymes, 1533
Galleria mellonella, 1533
penicillin, 1533
transaminases, 1533
mineral oil,
ethylene glycol, 723
nontarget insects, 723
preservatives, 723
propylene glycol, 723
mite, 586, 1928
allergy, 586
botanical insecticides, 586
ELISA, 1928
extraction, 1928
food, 586
grain, 586, 1928
storage, 1928
mixed viral infections,
fecundity, 646
host preference, 646
potato, 646

MOD system,
biological control, 283
insect rearing, 283
lacewing, 283
Verticel larval rearing unit, 283
molting hormone, 375, 1188
Chilo suppressalis, 375, 1188
imidacloprid, 375, 1188
juvenile hormone, 1188
juvenile hormone III, 375
rice plant, 375
monitoring, 343, 405, 1129, 1797
Bacillus thuringiensis, 1129
blacklight trap, 1797
Bt, 1129
European corn borer, 1129
lures, 343
Maconellicoccus hirsutus, 405
mesh cone trap, 1797
plum curculio, 343
release rate, 343
resistance management, 1129
sex pheromone, 405
sticky trap, 1797
temperature, 343
trapping, 405
water pan trap, 1797
monosultap,
Chilo suppressalis, 1854
fipronil, 1854
resistance, 1854
triazophos, 1854
mortality,
diatomaceous earth, 599
progeny production, 599
Prostephanus truncatus, 599
stored maize, 599
mortality rates,
biological control, 11
enzyme biomarkers, 11
pesticides, 11
mosquito,
larvicides, 61
Melanotaenia, 61
organophosphate, 61
sublethal effects, 61
moth,
attract and kill, 366
attractant, 366
behavior, 366
feeding, 366
movement, 54, 1037
breeding system, 1037
colony identification, 1037
Eoreuma loftini, 54
Mexican rice borer, 54
microsatellite genotyping, 1037
pheromone trap, 54
Rhinotermitidae, 1037
multicolored Asian lady beetle,
aggregation behavior, 1633
management, 1633
negative phototaxis, 1633
urban pest, 1633
multiple mating,
dispersal, 745
mass trapping, 745
pheromone, 745
suppression, 745
multiple resistance,
Brachiaria, 1896
competition, 1896
spittlebug, 1896
Musca domestica, 131, 155
Calliphora vicina, 131
germplasm regeneration, 131
house fly, 155
imidacloprid, 155
insect pollination, 131
leek, 131
methomyl, 155
spinosad, 155
mustard,
clone, 1764
okra, 1764
transmission efficiency, 1764
Mythimna separata,
AFLP, 1525
gene flow, 1525
genetic diversity, 1525
geographic population, 1525

N

NADPH-cytochrome P450 reductase,
Anopheles minimus, 946
CYP6AA3, 946
deltamethrin, 946
kinetics, 946
Nasonia,
Ecotilling, 1033
genome mapping, 1033
species identification, 1033
natural enemies,
introduced pest, 1268
Orius insidiosus, 1268
naturalyte,
biological control, 31
interaction, 31
Metarhizium, 31
wireworm, 31
navel orangeworm,
cold treatment, 1482
dried fruits, 1482
Indianmeal moth, 1482
tree nuts, 1482
near-infrared spectroscopy (NIRS),
insect fragments, 1714
internal insect infestation, 1714
rice weevil, 1714
stored products, 1714
negative cross-resistance,
Cyp12D1, 534
Cyp6g1, 534
cytochrome P450, 534
fumigant, 534
negative phototaxis,
aggregation behavior, 1633
management, 1633
multicolored Asian lady beetle, 1633
urban pest, 1633
neonicotinoid,
Popillia japonica, 431
sublethal effects, 431
toxicology, 431
neonicotinoids,
Anystis baccarum, 1210
carbamates, 1210
organophosphates, 1210
pyrethroids, 1210
Nezara viridula, 680, 1456
Acrosternum hilare, 1456
biocontrol, 680
Diabrotica, 680
Euschistus servus, 1456
host plant resistance, 1456
nonindigenous species,
ash utilization, 1577
eradication, 1577
Fraxinus, 1577
invasive pest, 1577
nonlinear models,
degree-days, 1098
Sancassania rodionovi, 1098
temperature-dependent development, 1098
nonrepellency,
delayed toxicity, 495
Reticulitermes hesperus, 495
termite movement, 495
nontarget insects,
ethylene glycol, 723
mineral oil, 723
preservatives, 723
propylene glycol, 723
nonwoven fabric,
adult beetle, 49
microbial control, 49
pine wilt disease, 49
novel application methods,
Chrysomela scripta, 1611
hybrid poplar, 1611
imidacloprid, 1611
nucleopolyhedrovirus,
fluorescent brightener, 1075
granulovirus, 1075
Mamestra brassicae, 1075
viral enhancement, 1075

nursery production,
Acer rubrum, 307
chlorpyrifos, 307
plant immersion, 307
Popillia japonica, 307
nutrition,
diet, 1062
rearing, 1062
root, 1062
weevil, 1062
nymphs,
Oebalus pugnax, 1282
Oryza sativa, 1282
rice panicle stage, 1282
rice stink bug, 1282
Nzi trap,
Greenhead trap, 613
Manitoba trap, 613
stable fly, 613
tabanids, 613

O

obliquebanded leafroller,
insecticide mixtures, 1831
insecticide resistance, 1831
integrated pest management, 1831
kaolin, 1831
Oebalus pugnax, 88, 1282
conservation tillage, 88
Helicoverpa zea, 88
nymphs, 1282
Oryza sativa, 1282
rice panicle stage, 1282
rice stink bug, 88, 1282
sorghum headworm, 88
okra,
clone, 1764
mustard, 1764
transmission efficiency, 1764
olive fly,
botanical insecticides, 1215
plant extracts, 1215
Rutaceae, 1215
onions,
emergence, 1194
overwintering, 1194
Thrips tabaci, 1194
volunteer onions, 1194
Opogona sacchari,
biosecurity, 1519
irradiation, 1519
quarantine treatment, 1519
sterilizing dose, 1519
orange oil extract,
Coptotermes formosanus, 880
d-limonene, 880
fumigation, 880
tunneling behavior, 880
organic agriculture,
aphid predators, 39
conservation biocontrol, 39
insectary crops, 39
Syrphidae, 39
organophosphate,
larvicides, 61
Melanotaenia, 61
mosquito, 61
sublethal effects, 61
organophosphates, 886, 1210
antagonism, 886
Anystis baccarum, 1210
Bemisia tabaci, 886
carbamates, 1210
neonicotinoids, 1210
potentiation, 886
pyrethroids, 886, 1210
oriental beetle,
Asiatic garden beetle, 1604
Bacillus thuringiensis serovar *japonensis*, 1604
European chafer, 1604
Japanese beetle, 1604
oriental fruit fly,
Bactrocera species, 670
classical biocontrol, 670
competitive displacement, 670
oriental fruit moth,
apple, 421
developmental rate, 421
peach, 421
population dynamics, 421
Orius insidiosus,
introduced pest, 1268
natural enemies, 1268
Oryza sativa,
nymphs, 1282
Oebalus pugnax, 1282
rice panicle stage, 1282
rice stink bug, 1282
Ostrinia nubilalis, 209, 1116, 1862
Bacillus thuringiensis, 1862
Bt, 1116
dispersal, 1116
European corn borer, 1116
proteinase-mediated resistance, 1862
resistance management, 1862
resistance mechanisms, 1862
Sesamia nonagrioides, 209
transgenic corn, 1116
Zea mays, 209
ovarian development,
female attractiveness, 1276
sex pheromone, 1276
Stenotus rubrovittatus, 1276
overwintering,
emergence, 1194
onions, 1194
Thrips tabaci, 1194
volunteer onions, 1194
ovicide,
insect growth regulator, 361
oxime ethers, 361
Spodoptera litura, 361
synergism, 361
oviposition substrate,
continuous in vitro culture, 26
hatching, 26
macrotype eggs, 26
oxime ethers,
insect growth regulator, 361
ovicide, 361
Spodoptera litura, 361
synergism, 361
Oxya chinensis,
acetylcholinesterase, 1409
general esterase, 1409
glutathione *S*-transferase, 1409
malathion, 1409

P

Paralobesia viteana,
Endopiza viteana, 855
insect growth regulator, 855
methoxyfenozide, 855
spinosad, 855
parasitoid-induced mortality,
Lariophagus distinguendus, 1011
life table, 1011
Sitophilus granarius, 1011
temperature, 1011
parasitoids,
Bemisia tabaci, 1053
Homalodisca lacerta, 1053
Homalodisca vitripennis, 1053
insecticide impacts, 1053
Paratrechina longicornis,
ant management, 509
bait preference, 509
crazy ants, 509
pathogenicity, 246, 291
Anastrepha ludens, 291
Bacillus thuringiensis, 246
Beauveria bassiana, 291
horizontal transmission, 291
Plutella xylostella, 246
septicemia, 246
virulence, 291
Xenorhabdus nematophila, 246
peach,
apple, 421
developmental rate, 421
oriental fruit moth, 421
population dynamics, 421
peanut,
Anticarsia gemmatalis, 1229
groundnut, 1229
insecticide efficacy, 1229
threecornered alfalfa hopper, 1229
tomato spotted wilt virus, 1229
pear,
apple, 702
CpGV, 702

ethyl (*E,Z*)-2,4-decadienoate, 702
kairomone, 702
Pelargonium peltatum,
integrated crop management, 1821
IPM, 1821
plant nutrition, 1821
Tetranychus urticae, 1821
Pelochrista womonana, 1248, 1555
Cylindrocopturus adspersus, 1248, 1555
Dectes texanus, 1248, 1555
pest management, 1248, 1555
sunflower, 1248, 1555
penicillin,
antioxidant enzymes, 1533
Galleria mellonella, 1533
midgut, 1533
transaminases, 1533
pepper,
aphids, 123
density thresholds, 123
tolerance, 123
peritrophic matrix,
Spodoptera litura, 20
Spodoptera litura nucleopolyhedrovirus, 20
Xestia c-nigrum granulovirus, 20
peritrophic membrane,
host plant resistance, 976
southwestern corn borer, 976
permethrin,
behavioral avoidance, 526
Plutella xylostella, 526
pyrethroids, 526
persistence,
abamectin, 411
azadirachtin, 411
Bemisia tabaci, 411
spinosad, 411
pest control,
abamectin susceptibility, 1637
Cacopsylla pyri, 1637
insecticide resistance, 1637
pest management, 459, 838, 1248, 1353, 1555, 1910
attraction, 1353
barriers, 1353
cabbage maggot, 838
Cylindrocopturus adspersus, 1248, 1555
Dectes texanus, 1248, 1555
diversification, 838
essential oils, 459
field sampling, 1910
fungus gnats, 459, 1353
greenhouse, 1353
intercropping, 838
leafminer, 1910
mealybugs, 459
Pelochrista womonana, 1248, 1555
relay cropping, 838
sunflower, 1248, 1555
Taylor's power law, 1910
urban green, 1910
western flower thrips, 459
pesticide metrics,
beneficials, 1741
IOBC, 1741
sustainable agriculture, 1741
pesticide residues,
methoxyfenozide, 773
reproductive parameters, 773
spinosad, 773
Spodoptera littoralis, 773
pesticides, 11, 1027
biocontrol, 1027
biological control, 11
enzyme biomarkers, 11
IOBC, 1027
mortality rates, 11
phenology, 847, 1104
emergence, 1104
greening disease, 847
huanglongbing, 847
kaolin, 847
maize, 1104
soybean, 1104
Surround, 847
pheromone, 745, 1815
dispersal, 745
grape berry moth, 1815
mass trapping, 745
mating disruption, 1815
multiple mating, 745
suppression, 745
pheromone dispenser,
emulsified wax, 1316
Grapholita molesta, 1316
mating disruption, 1316
pheromone trap, 54, 103
Eoreuma loftini, 54
feeding lure, 103
Korea, 103
Mexican rice borer, 54
movement, 54
Popillia japonica, 103
Popillia quadriguttata, 103
turfgrass, 103
pheromone trapping,
degree-days, 335
Heterocampa guttivitta, 335
population monitoring, 335
sugar maple, 335
photosynthesis,
gas exchange, 1692
host plant resistance, 1692
plant–insect interaction, 1692
Phyllocoptruta oleivora,
Aculops pelekassi, 233
presence–absence sampling, 233
proportions, 233
phytoplasma,
Bactericera cockerelli, 656
potato, 656
zebra chip, 656
phytosanitary treatment,
Agrilus planipennis, 1304
Fraxinus, 1304
invasive species, 1304
phytosanitary treatments,
fruit flies, 251
probit 9, 251
quarantine statistics, 251
PI 436684,
antibiosis, 1464
antixenosis, 1464
host plant resistance, 1464
PI 548664, 1464
PI 548664,
antibiosis, 1464
antixenosis, 1464
host plant resistance, 1464
PI 436684, 1464
PIC-Triad,
blueberry, 1596
imidacloprid, 1596
Japanese beetle, 1596
sublethal, 1596
pine wilt disease,
adult beetle, 49
microbial control, 49
nonwoven fabric, 49
Pisum sativum,
Bruchus pisorum, 1416
head capsule widths, 1416
transgenic peas, 1416
α-amylase inhibitor, 1416
plant extracts,
botanical insecticides, 1215
olive fly, 1215
Rutaceae, 1215
plant immersion,
Acer rubrum, 307
chlorpyrifos, 307
nursery production, 307
Popillia japonica, 307
plant nutrition,
integrated crop management, 1821
IPM, 1821
Pelargonium peltatum, 1821
Tetranychus urticae, 1821
plant resistance, 327, 566
bioassay, 566
community ecology, 327
Cry1Ab, 327
Diabrotica, 566
genetically modified, 327

Ipomoea, 566
maize, 327
plant–aphid interaction,
biotype, 990
Duraphis noxia, 990
elicitor, 990
Triticum aestivum, 990
plant–insect interaction,
gas exchange, 1692
host plant resistance, 1692
photosynthesis, 1692
plasticity,
developmental polymorphism, 627
instar number, 627
life history, 627
threshold size, 627
Platynota stultana,
carbon dioxide, 1084
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Pseudococcus maritimus, 1084
quarantine, 1084
plum curculio,
lures, 343
monitoring, 343
release rate, 343
temperature, 343
Plutella xylostella, 246, 526, 932
acetamiprid, 932
Bacillus thuringiensis, 246
behavioral avoidance, 526
inheritance, 932
pathogenicity, 246
permethrin, 526
pyrethroids, 526, 932
resistance, 932
septicemia, 246
Xenorhabdus nematophila, 246
pollination, 267, 1756
Apis mellifera, 267
Bombus terrestris, 1756
colony growth, 1756
drift, 1756
foraging behavior, 267
Vaccinium angustifolium, 267
polymerase chain reaction-restriction fragment length polymorphism,
Ceratitis capitata, 1444
mating success, 1444
sperm, 1444
spermathecae, 1444
Popillia japonica, 95, 103, 307, 431, 451, 830
Acer rubrum, 307
azadirachtin, 95, 451
biorational insecticides, 830
chlorpyrifos, 307
feeding lure, 103
feeding-induced aggregation, 95
hybrid tea rose, 830
imidacloprid, 95, 830
kaolin, 451
Korea, 103
limonene, 451
neonicotinoid, 431
nursery production, 307
pheromone traps, 103
plant immersion, 307
Popillia quadriguttata, 103
pyrethroid, 451
sublethal effects, 431
thresholds, 95
toxicology, 431
turfgrass, 103
Popillia quadriguttata,
feeding lure, 103
Korea, 103
pheromone traps, 103
Popillia japonica, 103
turfgrass, 103
population dynamics, 421, 1258
apple, 421
developmental rate, 421
invasive species, 1258
oriental fruit moth, 421
peach, 421
yield loss, 1258
population monitoring,
degree-days, 335
Heterocampa guttivitta, 335
pheromone trapping, 335
sugar maple, 335
portable EAG,
field studies, 1017
mating disruption, 1017
Pyralidae, 1017
stored-product pests, 1017
postharvest quality,
controlled atmosphere, 717
quarantine treatment, 717
ultralow oxygen, 717
western flower thrips, 717
postharvest treatment,
fruit flies, 1091
Litchi chinensis, 1091
quarantine pests, 1091
potato, 646, 656
Bactericera cockerelli, 656
fecundity, 646
host preference, 646
mixed viral infections, 646
phytoplasma, 656
zebra chip, 656
potentiation,
antagonism, 886
Bemisia tabaci, 886
organophosphates, 886
pyrethroids, 886
poultry litter,
Alphitobius diaperinus, 1136
darkling beetle, 1136
insect pest, 1136
Tenebrionidae, 1136
presence–absence sampling,
Aculops pelekassi, 233
Phyllocoptruta oleivora, 233
proportions, 233
preservatives,
ethylene glycol, 723
mineral oil, 723
nontarget insects, 723
propylene glycol, 723
Primetime,
fall armyworm, 1887
maysin, 1887
Zapalote Chico, 1887
probit 9,
fruit flies, 251
phytosanitary treatments, 251
quarantine statistics, 251
probit regression,
coverage probabilities, 1945
delta method, 1945
progeny production,
diatomaceous earth, 599
mortality, 599
Prostephanus truncatus, 599
stored maize, 599
projection pursuit regression,
counting, 225
Cucumis sativus, 225
Frankliniella occidentalis, 225
Rosa × *hybrida*, 225
proportions,
Aculops pelekassi, 233
Phyllocoptruta oleivora, 233
presence–absence sampling, 233
propylene glycol,
ethylene glycol, 723
mineral oil, 723
nontarget insects, 723
preservatives, 723
Prostephanus truncatus,
diatomaceous earth, 599
mortality, 599
progeny production, 599
stored maize, 599
protected stages,
Hyposoter didymator, 1789
insecticides, 1789
proteinase-mediated resistance,
Bacillus thuringiensis, 1862
Ostrinia nubilalis, 1862
resistance management, 1862
resistance mechanisms, 1862
Pseudococcus maritimus,
carbon dioxide, 1084
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
quarantine, 1084

Pseudoplusia includens,
Anticarsia gemmatalis, 962
host plant resistance, 962
Hypena scabra, 962
Pyralidae,
field studies, 1017
mating disruption, 1017
portable EAG, 1017
stored-product pests, 1017
pyrethroid, 451, 526, 886, 932, 1210
acetamiprid, 932
antagonism, 886
Anystis baccarum, 1210
azadirachtin, 451
behavioral avoidance, 526
Bemisia tabaci, 886
carbamates, 1210
inheritance, 932
kaolin, 451
limonene, 451
neonicotinoids, 1210
organophosphates, 886, 1210
permethrin, 526
Plutella xylostella, 526, 932
Popillia japonica, 451
potentiation, 886
resistance, 932
pyriproxyfen,
Bemisia tabaci, 1650
genetics, 1650
haplodiploid, 1650
insecticide resistance, 1650
Pyrrhalta luteola,
defoliation, 1808
Ulmus spp., 1808

Q

QTL,
fall armyworm, 1470
insect resistance, 1470
maize, 1470
southwestern corn borer, 1470
quantitative risk analysis,
Cylas formicarius elegantulus, 315
sweet potato, 315
quantitative trait loci, 1450, 1672
aphid, 1672
host resistance, 1672
Laodelphax striatellus, 1450
recombinant inbred line population, 1450
resistance, 1450
rice, 1450
Schizaphis graminum, 1672
simple sequence repeats, 1672
quarantine,
carbon dioxide, 1084
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
Pseudococcus maritimus, 1084
quarantine pests,
fruit flies, 1091
Litchi chinensis, 1091
postharvest treatment, 1091
quarantine statistics,
fruit flies, 251
phytosanitary treatments, 251
probit 9, 251
quarantine treatment, 717, 1499, 1519
biosecurity, 1519
codling moth, 1499
controlled atmosphere, 717
Cydia pomonella, 1499
irradiation, 1519
Opogona sacchari, 1519
postharvest quality, 717
Rhagoletis indifferens, 1499
sterilizing dose, 1519
ultralow oxygen, 717
western cherry fruit fly, 1499
western flower thrips, 717

R

R. flavipes,
C. formosanus, 475
hexaflumuron, 475
Sentricon, 475
raspberry ketone formate,
attractants, 1124
Bactrocera, 1124
melon fly, 1124
semiochemicals, 1124
rearing, 241, 1062
cellulose fiber, 241
diet, 1062
larval diet, 241
nutrition, 1062
root, 1062
screwworm, 241
spray-dried blood, 241
weevil, 1062
recombinant inbred line population,
Laodelphax striatellus, 1450
quantitative trait loci, 1450
resistance, 1450
rice, 1450
relative potencies,
canola, 1201
flea beetle, 1201
insecticides, 1201
lethal time, 1201
relay cropping,
cabbage maggot, 838
diversification, 838
intercropping, 838
pest management, 838
release rate,
lures, 343
monitoring, 343
plum curculio, 343
temperature, 343
reproduction,
development, 1476
life table parameters, 1476
Liposcelis tricolor, 1476
Wolbachia, 1476
reproductive parameters,
methoxyfenozide, 773
pesticide residues, 773
spinosad, 773
Spodoptera littoralis, 773
resampling software,
contaminant pest, 1000
Harmonia axyridis, 1000
wine grapes, 1000
resistance, 551, 932, 1450, 1587, 1854
acetamiprid, 932
apple orchard, 1587
azinphos-methyl, 551
Chilo suppressalis, 1854
codling moth, 1587
cross-resistance, 1587
Cydia pomonella, 551, 1587
fipronil, 1854
glutathione S-transferases, 551
inheritance, 932
insecticide, 551
Laodelphax striatellus, 1450
monosultap, 1854
Plutella xylostella, 932
pyrethroids, 932
quantitative trait loci, 1450
recombinant inbred line population, 1450
rice, 1450
triazophos, 1854
resistance management, 180, 1129, 1642, 1862
Bacillus thuringiensis, 180, 1129, 1642, 1862
Bt, 1129
cross-resistance, 180
European corn borer, 1129
Heliothis virescens, 180
high-dose/refuge strategy, 1642
mathematical modeling, 1642
monitoring, 1129
Ostrinia nubilalis, 1862
proteinase-mediated resistance, 1862
resistance mechanisms, 1862
RIDL, 1642
Vip3A, 180
resistance mechanisms,
Bacillus thuringiensis, 1862
Ostrinia nubilalis, 1862
proteinase-mediated resistance, 1862
resistance management, 1862

resistance monitoring, 164, 187
Bt maize, 164
cotton, 187
Diatraea saccharalis, 164
DNA diagnostics, 187
insecticide resistance management, 164
tobacco budworm, 187
retail stores,
augmentative biological control, 591
habitat complexity, 591
stored-products, 591
Reticulitermes flavipes,
soil barriers, 1391
termite baiting, 1391
Reticulitermes hesperus,
delayed toxicity, 495
nonrepellency, 495
termite movement, 495
reverse transcription-polymerase chain reaction,
enzyme-linked immunosorbent assay, 808
Fabaceae, 808
insect herbivory, 808
leaf area measurement, 808
Rhagoletis indifferens,
codling moth, 1499
Cydia pomonella, 1499
quarantine treatment, 1499
western cherry fruit fly, 1499
Rhinotermitidae,
breeding system, 1037
colony identification, 1037
microsatellite genotyping, 1037
movement, 1037
Rhyzopertha dominica, 604, 1724
diatomaceous earths, 1724
heat treatment, 604
laboratory bioassays, 1724
risk analysis, 604
Sitophilus oryzae, 604, 1724
Tribolium castaneum, 604
Tribolium confusum, 1724
ribosomal DNA,
COII mitochondrial DNA, 467
Coptotermes gestroi, 467
Coptotermes vastator, 467
synonymity, 467
rice,
Laodelphax striatellus, 1450
quantitative trait loci, 1450
recombinant inbred line population, 1450
resistance, 1450
rice panicle stage,
nymphs, 1282
Oebalus pugnax, 1282
Oryza sativa, 1282
rice stink bug, 1282
rice plant,
Chilo suppressalis, 375
imidacloprid, 375
juvenile hormone III, 375
molting hormone, 375
rice stink bug, 88, 1282
conservation tillage, 88
Helicoverpa zea, 88
nymphs, 1282
Oebalus pugnax, 88, 1282
Oryza sativa, 1282
rice panicle stage, 1282
sorghum headworm, 88
rice weevil,
insect fragments, 1714
internal insect infestation, 1714
near-infrared spectroscopy (NIRS), 1714
stored products, 1714
RIDL,
Bacillus thuringiensis, 1642
high-dose/refuge strategy, 1642
mathematical modeling, 1642
resistance management, 1642
risk analysis,
heat treatment, 604
Rhyzopertha dominica, 604
Sitophilus oryzae, 604
Tribolium castaneum, 604
root,
diet, 1062
nutrition, 1062
rearing, 1062
weevil, 1062
Rosa × *hybrida*,
counting, 225
Cucumis sativus, 225
Frankliniella occidentalis, 225
projection pursuit regression, 225
Russian stock,
honey bees, 258
integrative pest management, 258
tracheal mite, 258
varroa mite, 258
Rutaceae,
botanical insecticides, 1215
olive fly, 1215
plant extracts, 1215

S

S. chacoense-derived resistance,
Bacillus thuringiensis (Bt), 573
cry1Ia1, 573
cry3A, 573
glandular trichomes, 573
SADIE, 790, 1773
cultural management, 1773
Ectomyelois ceratoniae, 1773
geographic information system, 1773
geospatial mapping, 790
geostatistics, 790
soil insects, 790
sound, 790
sampling, 136, 1160, 1622
Bemisia tabaci, 1160
beneficial arthropods, 136
Blissus occiduus, 136
blueberries, 1622
buffalograss, 136
cucumber, 1160
distribution, 1622
hot-spots, 1622
thrips, 1622
western chinch bugs, 136
within-plant distribution, 1160
Sancassania rodionovi,
degree-days, 1098
nonlinear models, 1098
temperature-dependent development, 1098
Saperda vestita,
chipping, 1328
systemic insecticides, 1328
Tilia spp., 1328
sawfly,
bread wheat, 79
Cephus cinctus, 79
economic loss, 79
yield loss, 79
Schizaphis graminum,
aphid, 1672
host resistance, 1672
quantitative trait loci, 1672
simple sequence repeats, 1672
Scolytidae,
Cleridae, 815
Conophthorus coniperda, 815
Enoclerus nigripes, 815
kairomone, 815
screwworm,
cellulose fiber, 241
larval diet, 241
rearing, 241
spray-dried blood, 241
seasonal ecology,
golf course, 1344
Listronotus maculicollis, 1344
turfgrass insect, 1344
seed-treatment,
canola, 765
clothianidin, 765
honey bees, 765
long-term assessment, 765
semiochemical baits,
Diabrotica, 1542
IPM, 1542

landscape, 1542
semiochemicals,
attractants, 1124
Bactrocera, 1124
melon fly, 1124
raspberry ketone formate, 1124
Sentricon,
C. formosanus, 475
hexaflumuron, 475
R. flavipes, 475
septicemia,
Bacillus thuringiensis, 246
pathogenicity, 246
Plutella xylostella, 246
Xenorhabdus nematophila, 246
Sesamia nonagrioides, 195, 209
Bacillus thuringiensis, 195
F_2 screen resistance management, 195
high-dose/refuge strategy, 195
Ostrinia nubilalis, 209
transgenic maize, 195
Zea mays, 209
sex pheromone, 405, 1276
female attractiveness, 1276
Maconellicoccus hirsutus, 405
monitoring, 405
ovarian development, 1276
Stenotus rubrovittatus, 1276
trapping, 405
sexual behavior,
Grapholita molesta, 72
methoxyfenozide, 72
wind tunnel, 72
sharka,
herbaceous species, 1047
holocycle, 1047
vector, 1047
weeds, 1047
silk-feeding bioassay,
Helicoverpa zea, 1704
husk extension, 1704
Sitophilus zeamais, 1704
stink bugs, 1704
silkworm,
amplified fragment length polymorphism markers, 939
Bombyx mori, 939
genetic diversity, 939
simple sequence repeats,
aphid, 1672
host resistance, 1672
quantitative trait loci, 1672
Schizaphis graminum, 1672
Sipha maydis,
cereal pest, 1781
geographical distribution, 1781
host preference, 1781
site-specific management,
corn, 781
management zones, 781
Sitophilus granarius,
Lariophagus distinguendus, 1011
life table, 1011
parasitoid-induced mortality, 1011
temperature, 1011
Sitophilus oryzae, 604, 1724
diatomaceous earths, 1724
heat treatment, 604
laboratory bioassays, 1724
Rhyzopertha dominica, 604, 1724
risk analysis, 604
Tribolium castaneum, 604
Tribolium confusum, 1724
Sitophilus zeamais,
Helicoverpa zea, 1704
husk extension, 1704
silk-feeding bioassay, 1704
stink bugs, 1704
soil barriers,
Reticulitermes flavipes, 1391
termite baiting, 1391
soil insects,
geospatial mapping, 790
geostatistics, 790
SADIE, 790
sound, 790
Solanum tuberosum,
Bacillus thuringiensis, 172
genetic transformation, 172
insect resistance, 172
Tecia solanivora, 172
Solbait,
Anastrepha ludens, 440
GF-120, 440
Mexican fruit fly, 440
soldier proportions,
colony density, 1381
colony size, 1381
cuticular hydrocarbons, 1381
mark–release–recapture, 1381
Solenopotes capillatus,
Anoplura, 619
little blue cattle louse, 619
Solenopsis invicta,
competition, 148
interspecific aggression, 148
Linepithema humile, 148
sorghum headworm,
conservation tillage, 88
Helicoverpa zea, 88
Oebalus pugnax, 88
rice stink bug, 88
sound,
geospatial mapping, 790
geostatistics, 790
SADIE, 790
soil insects, 790
southern green stink bug,
cotton, 1560
heat units, 1560
injury levels, 1560
integrated pest management, 1560
southern pine beetle,
geographic information system, 381
hazard rating, 381
Landsat ETM+, 381
southwestern corn borer, 969, 976, 1470
fall armyworm, 1470
host plant resistance, 969, 976
insect resistance, 1470
maize, 1470
peritrophic membrane, 976
QTL, 1470
soybean,
emergence, 1104
maize, 1104
phenology, 1104
species identification, 759, 1033
Ecotilling, 1033
genome mapping, 1033
Helicoverpa zea, 759
Heliothis virescens, 759
Nasonia, 1033
sperm,
Ceratitis capitata, 1444
mating success, 1444
polymerase chain reaction-restriction fragment length polymorphism, 1444
spermathecae, 1444
spermathecae,
Ceratitis capitata, 1444
mating success, 1444
polymerase chain reaction-restriction fragment length polymorphism, 1444
sperm, 1444
spider mites,
bedding plants, 1904
damage threshold, 1904
esthetic damage, 1904
floriculture, 1904
spinosad, 155, 411, 773, 855
abamectin, 411
azadirachtin, 411
Bemisia tabaci, 411
Endopiza viteana, 855
house fly, 155
imidacloprid, 155
insect growth regulator, 855
methomyl, 155
methoxyfenozide, 773, 855
Musca domestica, 155

Paralobesia viteana, 855
persistence, 411
pesticide residues, 773
reproductive parameters, 773
Spodoptera littoralis, 773
spittlebug,
Brachiaria, 1896
competition, 1896
multiple resistance, 1896
Spodoptera frugiperda, 954, 1569
armyworm, 954
cotton, 1569
cytochrome oxidase I, 954
fall armyworm, 1569
host strains, 1569
stable isotope, 1569
Spodoptera littoralis,
methoxyfenozide, 773
pesticide residues, 773
reproductive parameters, 773
spinosad, 773
Spodoptera litura, 20, 361
insect growth regulator, 361
ovicide, 361
oxime ethers, 361
peritrophic matrix, 20
Spodoptera litura nucleopolyhedrovirus, 20
synergism, 361
Xestia c-nigrum granulovirus, 20
Spodoptera litura nucleopolyhedrovirus,
peritrophic matrix, 20
Spodoptera litura, 20
Xestia c-nigrum granulovirus, 20
spray-dried blood,
cellulose fiber, 241
larval diet, 241
rearing, 241
screwworm, 241
sprayable microencapsulated pheromone,
mating disruption, 1360
Tortricidae, 1360
stable fly,
Greenhead trap, 613
Manitoba trap, 613
Nzi trap, 613
tabanids, 613
stable isotope,
cotton, 1569
fall armyworm, 1569
host strains, 1569
Spodoptera frugiperda, 1569
Steinernema,
blackberries, 398
entomopathogenic nematodes, 398
Heterorhabditis, 398
insecticides, 398
Stenotus rubrovittatus,
female attractiveness, 1276
ovarian development, 1276
sex pheromone, 1276
sterile insect technique, 273, 730, 752, 1153, 1173, 1180
Anastrepha ludens, 752, 1153
Anastrepha serpentina, 1173
aromatherapy, 273
calling behavior, 1173
Ceratitis capitata, 1180
Cylas formicarius elegantulus, 730
dispersal ability, 730
gamma radiation, 730
irradiation doses, 1153
male age, 1180
mass rearing, 1173
mating competitiveness, 752, 1153
mating success, 1180
Mediterranean fruit fly, 273, 1180
sterility induction, 1153
strain replacement, 752
sweetpotato weevil, 730
sterility induction,
Anastrepha ludens, 1153
irradiation doses, 1153
mating competitiveness, 1153
sterile insect technique, 1153
sterilizing dose,
biosecurity, 1519
irradiation, 1519
Opogona sacchari, 1519
quarantine treatment, 1519
sticky trap,
blacklight trap, 1797
mesh cone trap, 1797
monitoring, 1797
water pan trap, 1797
stink bugs,
Helicoverpa zea, 1704
husk extension, 1704
silk-feeding bioassay, 1704
Sitophilus zeamais, 1704
storage,
ELISA, 1928
extraction, 1928
grain, 1928
mite, 1928
stored maize,
diatomaceous earth, 599
mortality, 599
progeny production, 599
Prostephanus truncatus, 599
stored-product pests,
field studies, 1017
mating disruption, 1017
portable EAG, 1017
Pyralidae, 1017
stored-products, 591, 1714, 1924
aggregation pheromone, 1924
augmentative biological control, 591
dimethyldecanal, 1924
habitat complexity, 591
insect fragments, 1714
internal insect infestation, 1714
near-infrared spectroscopy (NIRS), 1714
retail stores, 591
rice weevil, 1714
trap response, 1924
strain replacement,
Anastrepha ludens, 752
mating competitiveness, 752
sterile insect technique, 752
sublethal, 1512, 1596
blueberry, 1596
emergence inhibition, 1512
imidacloprid, 1596
insecticides, 1512
Japanese beetle, 1596
PIC-Triad, 1596
survival, 1512
sublethal effects, 61, 431
larvicides, 61
Melanotaenia, 61
mosquito, 61
neonicotinoid, 431
organophosphate, 61
Popillia japonica, 431
toxicology, 431
sugar maple,
degree-days, 335
Heterocampa guttivitta, 335
pheromone trapping, 335
population monitoring, 335
sunflower, 1248, 1555
Cylindrocopturus adspersus, 1248, 1555
Dectes texanus, 1248, 1555
Pelochrista womonana, 1248, 1555
pest management, 1248, 1555
suppression,
dispersal, 745
mass trapping, 745
multiple mating, 745
pheromone, 745
surfactant,
Blattella germanica, 871
synergism, 871
Surround,
greening disease, 847
huanglongbing, 847
kaolin, 847
phenology, 847
survival,
emergence inhibition, 1512
insecticides, 1512

sublethal, 1512
sustainable agriculture,
beneficials, 1741
IOBC, 1741
pesticide metrics, 1741
swarming behaviors,
alate monitoring, 1400
Coptotermes formosanus, 1400
invasive species, 1400
termite survey, 1400
sweet potato,
Cylas formicarius elegantulus, 315
quantitative risk analysis, 315
sweetpotato weevil,
Cylas formicarius elegantulus, 730
dispersal ability, 730
gamma radiation, 730
sterile insect technique, 730
sweetpotato whitefly,
cotton aphid, 1428
cotton bollworm, 1428
cotton fleahopper, 1428
western flower thrips, 1428
synergism, 361, 871
Blattella germanica, 871
insect growth regulator, 361
ovicide, 361
oxime ethers, 361
Spodoptera litura, 361
surfactant, 871
synergists,
acetylcholinesterase, 1836
citrus, 1836
esterases, 1836
insecticides, 1836
synonymity,
COII mitochondrial DNA, 467
Coptotermes gestroi, 467
Coptotermes vastator, 467
ribosomal DNA, 467
Syrphidae,
aphid predators, 39
conservation biocontrol, 39
insectary crops, 39
organic agriculture, 39
systemic insecticides,
chipping, 1328
Saperda vestita, 1328
Tilia spp., 1328

T

tabanids,
Greenhead trap, 613
Manitoba trap, 613
Nzi trap, 613
stable fly, 613
Taylor's power law,
field sampling, 1910
leafminer, 1910
pest management, 1910
urban green, 1910
Tecia solanivora,
Bacillus thuringiensis, 172
genetic transformation, 172
insect resistance, 172
Solanum tuberosum, 172
temperature, 343, 357, 1011
Earias vitella, 357
insecticide toxicity, 357
Lariophagus distinguendus, 1011
life table, 1011
lures, 343
monitoring, 343
parasitoid-induced mortality, 1011
plum curculio, 343
release rate, 343
Sitophilus granarius, 1011
temperature dependent,
cucumber, 350
degree-days, 350
development, 350
Liriomyza sativae, 350
temperature-dependent development,
degree-days, 1098
nonlinear models, 1098
Sancassania rodionovi, 1098
Tenebrionidae,
Alphitobius diaperinus, 1136
darkling beetle, 1136
insect pest, 1136
poultry litter, 1136
Tephritidae,
ammonia quantification, 580
Anastrepha ludens, 580
Ceratitis capitata, 580
Fourier transform spectroscopy, 580
termite,
termite attack and damage, 488
termiticide efficacy testing, 488
termite attack and damage,
termite, 488
termiticide efficacy testing, 488
termite baiting,
Reticulitermes flavipes, 1391
soil barriers, 1391
termite movement,
delayed toxicity, 495
nonrepellency, 495
Reticulitermes hesperus, 495
termite survey,
alate monitoring, 1400
Coptotermes formosanus, 1400
invasive species, 1400
swarming behaviors, 1400
termiticide efficacy testing,
termite, 488
termite attack and damage, 488
Tetranychus cinnabarinus,
acaricidal activity, 1298
Juglans regia, 1298
Tetranychus viennensis, 1298
Tetranychus pacificus,
carbon dioxide, 1084
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
Pseudococcus maritimus, 1084
quarantine, 1084
Tetranychus urticae,
integrated crop management, 1821
IPM, 1821
Pelargonium peltatum, 1821
plant nutrition, 1821
Tetranychus viennensis,
acaricidal activity, 1298
Juglans regia, 1298
Tetranychus cinnabarinus, 1298
Thailand,
Aedes (*Stegomyia*), 545
diagnostic dose, 545
insecticide, 545
'Thompson Seedless',
carbon dioxide, 1084
controlled atmosphere, 1084
Frankliniella occidentalis, 1084
Platynota stultana, 1084
Pseudococcus maritimus, 1084
quarantine, 1084
threecornered alfalfa hopper,
Anticarsia gemmatalis, 1229
groundnut, 1229
insecticide efficacy, 1229
peanut, 1229
tomato spotted wilt virus, 1229
threshold size,
developmental polymorphism, 627
instar number, 627
life history, 627
plasticity, 627
thresholds,
azadirachtin, 95
feeding-induced aggregation, 95
imidacloprid, 95
Popillia japonica, 95
thrips,
blueberries, 1622
distribution, 1622
hot-spots, 1622
sampling, 1622
Thrips tabaci,
emergence, 1194
onions, 1194

overwintering, 1194
volunteer onions, 1194
Thuja,
exotic beetle, 1291
host preference, 1291
Juniperus, 1291
trap log, 1291
Thysanoptera,
inheritance, 916
insecticide resistance, 916
Tilia spp.,
chipping, 1328
Saperda vestita, 1328
systemic insecticides, 1328
tobacco budworm,
cotton, 187
DNA diagnostics, 187
resistance monitoring, 187
tobacco thrips,
advisory, 1241
integrated pest management, 1241
tomato spotted wilt virus, 1241
virginia market-type peanut, 1241
tolerance, 123, 215
antibiosis, 215
antixenosis, 215
aphids, 123
crucifers, 215
density thresholds, 123
diamondback moth, 215
pepper, 123
tomato spotted wilt virus, 1229, 1241
advisory, 1241
Anticarsia gemmatalis, 1229
groundnut, 1229
insecticide efficacy, 1229
integrated pest management, 1241
peanut, 1229
threecornered alfalfa hopper, 1229
tobacco thrips, 1241
virginia market-type peanut, 1241
Tortricidae,
mating disruption, 1360
sprayable microencapsulated pheromone, 1360
toxicant,
attractant, 483
bait, 483
house fly, 483
trap, 483
toxicology,
neonicotinoid, 431
Popillia japonica, 431
sublethal effects, 431
tracheal mite,
honey bees, 258
integrative pest management, 258
Russian stock, 258
varroa mite, 258
transaminases,
antioxidant enzymes, 1533
Galleria mellonella, 1533
midgut, 1533
penicillin, 1533
transgenic corn, 1116, 1880
Bt, 1116
Bt resistance management, 1880
Cry1Ab protein, 1880
Diatraea saccharalis, 1880
dispersal, 1116
European corn borer, 1116
Ostrinia nubilalis, 1116
transgenic maize,
Bacillus thuringiensis, 195
F_2 screen resistance management, 195
high-dose/refuge strategy, 195
Sesamia nonagrioides, 195
transgenic peas,
Bruchus pisorum, 1416
head capsule widths, 1416
Pisum sativum, 1416
α-amylase inhibitor, 1416
transmission efficiency,
clone, 1764
mustard, 1764
okra, 1764
trap, 483, 737, 1147
AFF lure, 1147
Anastrepha ludens, 1147
apple, 737
attractant, 483, 1147
bait, 483
BioLure, 1147
Conotrachelus nenuphar, 737
highbush blueberry, 737
house fly, 483
toxicant, 483
trap log,
exotic beetle, 1291
host preference, 1291
Juniperus, 1291
Thuja, 1291
trap response,
aggregation pheromone, 1924
dimethyldecanal, 1924
stored products, 1924
trapping,
Maconellicoccus hirsutus, 405
monitoring, 405
sex pheromone, 405
treatment threshold,
damage relationships, 801
flue-cured tobacco, 801
infestation, 801
tree nuts,
cold treatment, 1482
dried fruits, 1482
Indianmeal moth, 1482
navel orangeworm, 1482
Trialeurodes vaporariorum,
light-diffusion, 389
UV-blocking films, 389
whitefly, 389
triazophos,
Chilo suppressalis, 1854
fipronil, 1854
monosultap, 1854
resistance, 1854
Tribolium castaneum, 604, 1071
Beauveria bassiana, 1071
desiccation, 1071
flour beetle, 1071
heat treatment, 604
humidity, 1071
Rhyzopertha dominica, 604
risk analysis, 604
Sitophilus oryzae, 604
Tribolium confusum,
diatomaceous earths, 1724
laboratory bioassays, 1724
Rhyzopertha dominica, 1724
Sitophilus oryzae, 1724
Triticum aestivum, 990, 1685
biotype, 990, 1685
Diuraphis noxia, 1685
Duraphis noxia, 990
elicitor, 990
Hordeum vulgare, 1685
host plant resistance, 1685
plant–aphid interaction, 990
tunneling behavior,
Coptotermes formosanus, 880
d-limonene, 880
fumigation, 880
orange oil extract, 880
turfgrass,
feeding lure, 103
Korea, 103
pheromone traps, 103
Popillia japonica, 103
Popillia quadriguttata, 103
turfgrass insect,
golf course, 1344
Listronotus maculicollis, 1344
seasonal ecology, 1344

U

Ulmus spp.,
defoliation, 1808
Pyrrhalta luteola, 1808
ultralow oxygen,
controlled atmosphere, 717
postharvest quality, 717
quarantine treatment, 717
western flower thrips, 717

urban green,
field sampling, 1910
leafminer, 1910
pest management, 1910
Taylor's power law, 1910
urban pest,
aggregation behavior, 1633
management, 1633
multicolored Asian lady beetle, 1633
negative phototaxis, 1633
UV-blocking films, 110, 389
diffusion, 110
dispersal, 110
Encarsia formosa, 110
light-diffusion, 389
Trialeurodes vaporariorum, 389
whitefly, 389

V

Vaccinium angustifolium,
Apis mellifera, 267
foraging behavior, 267
pollination, 267
Varroa destructor,
Apis mellifera, 1
Beauveria bassiana, 1
hive treatment, 1
varroa mite,
honey bees, 258
integrative pest management, 258
Russian stock, 258
tracheal mite, 258
vector,
herbaceous species, 1047
holocycle, 1047
sharka, 1047
weeds, 1047
vector transmission,
Chrysanthemum yellows phytoplasma, 1504
Euscelidius variegatus, 1504
Euscelis incisus, 1504
Macrosteles quadripunctulatus, 1504
Verticel larval rearing unit,
biological control, 283
insect rearing, 283
lacewing, 283
MOD system, 283
Vip3A,
Bacillus thuringiensis, 180
cross-resistance, 180
Heliothis virescens, 180
resistance management, 180
viral enhancement,
fluorescent brightener, 1075
granulovirus, 1075
Mamestra brassicae, 1075
nucleopolyhedrovirus, 1075
virginia market-type peanut,
advisory, 1241
integrated pest management, 1241
tobacco thrips, 1241
tomato spotted wilt virus, 1241
virulence,
Anastrepha ludens, 291
Beauveria bassiana, 291
horizontal transmission, 291
pathogenicity, 291
volunteer onions,
emergence, 1194
onions, 1194
overwintering, 1194
Thrips tabaci, 1194

W

water pan trap,
blacklight trap, 1797
mesh cone trap, 1797
monitoring, 1797
sticky trap, 1797
weeds, 298, 1047
Agromyzidae, 298
conservation biological control, 298
ecological infrastructures, 298
herbaceous species, 1047
holocycle, 1047
hymenopteran parasitoids, 298
sharka, 1047
vector, 1047
weevil,
diet, 1062
nutrition, 1062
rearing, 1062
root, 1062
western cherry fruit fly,
codling moth, 1499
Cydia pomonella, 1499
quarantine treatment, 1499
Rhagoletis indifferens, 1499
western chinch bugs,
beneficial arthropods, 136
Blissus occiduus, 136
buffalograss, 136
sampling, 136
western flower thrips, 459, 717, 1428
controlled atmosphere, 717
cotton aphid, 1428
cotton bollworm, 1428
cotton fleahopper, 1428
essential oils, 459
fungus gnats, 459
mealybugs, 459
pest management, 459
postharvest quality, 717
quarantine treatment, 717
sweetpotato whitefly, 1428
ultralow oxygen, 717
wheat,
barley, 1679
host plant resistance, 1679
host race, 1679
wheat germ oil,
Bactrocera dorsalis, 322
fruit fly rearing, 322
liquid diet, 322
whitefly,
light-diffusion, 389
Trialeurodes vaporariorum, 389
UV-blocking films, 389
wind tunnel,
Grapholita molesta, 72
methoxyfenozide, 72
sexual behavior, 72
wine grapes,
contaminant pest, 1000
Harmonia axyridis, 1000
resampling software, 1000
wireworm,
biological control, 31
interaction, 31
Metarhizium, 31
naturalyte, 31
within-plant distribution,
Bemisia tabaci, 1160
cucumber, 1160
sampling, 1160
Wolbachia,
development, 1476
life table parameters, 1476
Liposcelis tricolor, 1476
reproduction, 1476
wood protectants,
botanic termiticide, 866
Ku Shen, 866

X

Xenopsylla cheopis,
Aedes aegypti, 622
biocidal, 622
essential oils, 622
Ixodes scapularis, 622
Xenorhabdus nematophila,
Bacillus thuringiensis, 246
pathogenicity, 246
Plutella xylostella, 246
septicemia, 246
Xestia c-nigrum granulovirus,
peritrophic matrix, 20
Spodoptera litura, 20
Spodoptera litura nucleopolyhedrovirus, 20

Y

yellow sticky trap,
Acalymma vittatum, 823
Cucumis melo, 823
Diabrotica undecimpunctata howardi, 823

direct scouting, 823
yield,
canola, 710
flea beetle, 710
injury, 710
insecticides, 710
yield loss, 79, 1258
bread wheat, 79
Cephus cinctus, 79
economic loss, 79
invasive species, 1258
population dynamics, 1258
sawfly, 79

Z

(Z)-9-tricosene,
behavioral resistance, 1489
fly bait, 1489
house flies, 1489
methomyl, 1489
Zapalote Chico,
fall armyworm, 1887
maysin, 1887
Primetime, 1887
Zea mays,
Ostrinia nubilalis, 209
Sesamia nonagrioides, 209
zebra chip,
Bactericera cockerelli, 656
phytoplasma, 656
potato, 656